

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : A61K 38/00, C07K 5/00, 7/00, 16/00, 17/00, C07H 21/04, C12N 15/00, 15/09, 15/63, 15/70, 15/74, C12P 21/06, C12Q 1/68</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/42120</p>
<p>(22) International Filing Date: 18 February 1999 (18.02.99)</p>		<p>(43) International Publication Date: 26 August 1999 (26.08.99)</p>
<p>(30) Priority Data: 09/027,337 20 February 1998 (20.02.98) US</p>		<p>(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b> <i>With international search report.</i></p>
<p>(71) Applicant: THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ARKANSAS [US/US]; 2404 North University Avenue, Little Rock, AR 72207-3608 (US).</p>		
<p>(72) Inventors: O'BRIEN, Timothy, J.; 2625 Grist Mill Road, Little Rock, AR 72227 (US). TANIMOTO, Hirotochi; Apartment 804, 701 Green Mountain Drive, Little Rock, AR 72211 (US).</p>		
<p>(74) Agent: ADLER, Benjamin, A.; McGregor &amp; Adler, 8011 Candle Lane, Houston, TX 77071 (US).</p>		
<p>(54) Title: TADG-15: AN EXTRACELLULAR SERINE PROTEASE OVEREXPRESSED IN BREAST AND OVARIAN CARCINOMAS</p>		
<p>(57) Abstract</p> <p>The present invention provides a DNA encoding a TADG-15 protein selected from the group consisting of: (a) isolated DNA which encodes a TADG-15 protein; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a TADG-15 protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to the degeneracy of the genetic code, and which encodes a TADG-15 protein. Also provided is a vector capable of expressing the DNA of the present invention adapted for expression in a recombinant cell and regulatory elements necessary for expression of the DNA in the cell.</p>		

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

5           TADG-15: AN EXTRACELLULAR SERINE PROTEASE  
OVEREXPRESSED IN BREAST AND OVARIAN CARCINOMAS

10

BACKGROUND OF THE INVENTION

15   Field of the Invention

The present invention relates generally to the fields of cellular biology and the diagnosis of neoplastic disease. More specifically, the present invention relates to an extracellular serine protease termed Tumor Antigen Derived Gene-15 (TADG-15), which is  
20 overexpressed in breast and ovarian carcinomas.

Description of the Related Art

Extracellular proteases have been directly associated with tumor growth, shedding of tumor cells and invasion of target organs.  
25 Individual classes of proteases are involved in, but not limited to (1) the digestion of stroma surrounding the initial tumor area, (2) the digestion of the cellular adhesion molecules to allow dissociation of tumor cells; and (3) the invasion of the basement membrane for

metastatic growth and the activation of both tumor growth factors and angiogenic factors.

The prior art is deficient in the lack of effective means of screening to identify proteases overexpressed in carcinoma. The  
5 present invention fulfills this longstanding need and desire in the art.

### SUMMARY OF THE INVENTION

10 The present invention discloses a screening program to identify proteases overexpressed in carcinoma by examining PCR products amplified using differential display in early stage tumors, metastatic tumors compared to that of normal tissues.

In one embodiment of the present invention, there is  
15 provided a DNA encoding a TADG-15 protein selected from the group consisting of: (a) isolated DNA which encodes a TADG-15 protein; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a TADG-15 protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to the  
20 degeneracy of the genetic code, and which encodes a TADG-15 protein.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention adapted for expression in a recombinant cell and regulatory  
25 elements necessary for expression of the DNA in the cell.

In yet another embodiment of the present invention, there is provided a host cell transfected with the vector of the present invention, the vector expressing a TADG-15 protein.

In still yet another embodiment of the present invention, there is provided a method of detecting expression of a TADG-15 mRNA, comprising the steps of: (a) contacting mRNA obtained from the cell with the labeled hybridization probe; and (b) detecting  
5 hybridization of the probe with the mRNA.

Other and further aspects, features, and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention given for the purpose of disclosure.

10

### BRIEF DESCRIPTION OF THE DRAWINGS

So that the matter in which the above-recited features, advantages and objects of the invention, as well as others which will  
15 become clear, are attained and can be understood in detail, more particular descriptions of the invention briefly summarized above may be had by reference to certain embodiments thereof which are illustrated in the appended drawings. These drawings form a part of the specification. It is to be noted, however, that the appended  
20 drawings illustrate preferred embodiments of the invention and therefore are not to be considered limiting in their scope.

**Figure 1** shows a comparison of PCR products derived from normal and breast carcinoma cDNA as shown by staining in an agarose gel.

25 **Figure 2** shows a comparison of the serine protease catalytic domain of TADG-15 (SEQ ID No: 14) with hepsin (Heps, SEQ ID No: 3), (Scce, SEQ ID No: 4), trypsin (Try, SEQ ID No: 5), chymotrypsin (Chymb, SEQ ID No: 6), factor 7 (Fac7, SEQ ID No:

7) and tissue plasminogen activator (Tpa, SEQ ID No: 8). The asterisks indicate conserved amino acids of catalytic triad.

Figure 3 shows quantitative PCR analysis of TADG-15 expression.

5 Figure 4 shows the ratio of TADG-15 expression to expression of  $\beta$ -tubulin in normal tissues, low malignant potential tumors (LMP) and carcinomas.

Figure 5 shows the TADG-15 expression in tumor cell lines derived from both ovarian and breast carcinoma tissues.

10 Figure 6 shows the overexpression of TADG-15 in other tumor tissues.

Figure 7 shows the Northern blots of TADG-15 expression in ovarian carcinomas, fetal and normal adult tissues.

15 Figure 8 shows a diagram of the TADG-15 transcript and the clones with the origin of their derivation.

Figure 9 shows nucleotide sequence of the TADG-15 cDNA (SEQ ID No: 1) and amino acid sequence of the TADG-15 protein (SEQ ID No: 2)

20 Figure 10 shows the amino acid sequence of the TADG-15 protease including functional sites and domains.

Figure 11 shows a structure diagram of the TADG-15 protein including functional domains.

Figure 12 shows a nucleotide sequence comparison between TADG-15 and human SNC-19 (GeneBank accession #U20428).

25

## DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "cDNA" shall refer to the DNA copy of the mRNA transcript of a gene.

As used herein, the term "derived amino acid sequence" shall mean the amino acid sequence determined by reading the triplet  
5 sequence of nucleotide bases in the cDNA.

As used herein the term "screening a library" shall refer to the process of using a labeled probe to check whether, under the appropriate conditions, there is a sequence complementary to the probe present in a particular DNA library. In addition, "screening a  
10 library" could be performed by PCR.

As used herein, the term "PCR" refers to the polymerase chain reaction that is the subject of U.S. Patent Nos. 4,683,195 and 4,683,202 to Mullis, as well as other improvements now known in the art.

15 The TADG-15 cDNA is 3147 base pairs long (SEQ ID No:1) and encoding for a 855 amino acid protein (SEQ ID No:2). The availability of the TADG-15 gene opens the way for a number studies that can lead to various applications. For example, the TADG-15 gene can be used as a diagnostic or therapeutic target in ovarian carcinoma  
20 and other carcinomas including breast, prostate, lung and colon.

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis,  
25 Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" [B.D. Hames & S.J. Higgins eds. (1985)];

"Transcription and Translation" [B.D. Hames & S.J. Higgins eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

5           Therefore, if appearing herein, the following terms shall have the definitions set out below.

          The amino acid described herein are preferred to be in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired  
10 functional property of immunoglobulin-binding is retained by the polypeptide. NH<sub>2</sub> refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 243:3552-59  
15 (1969), abbreviations for amino acid residues are shown in the following Table of Correspondence:



TABLE OF CORRESPONDENCE

	<u>SYMBOL</u> <u>1-Letter</u>	<u>3-Letter</u>	<u>AMINO ACID</u>
5	Y	Tyr	tyrosine
	G	Gly	glycine
	F	Phe	Phenylalanine
	M	Met	methionine
	A	Ala	alanine
10	S	Ser	serine
	I	Ile	isoleucine
	L	Leu	leucine
	T	Thr	threonine
	V	Val	valine
15	P	Pro	proline
	K	Lys	lysine
	H	His	histidine
	Q	Gln	glutamine
	E	Glu	glutamic acid
20	W	Trp	tryptophan
	R	Arg	arginine
	D	Asp	aspartic acid
	N	Asn	asparagine
	C	Cys	cysteine
25			

It should be noted that all amino-acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino-acid residues. The above Table is presented to correlate the three-letter and one-letter notations which may appear alternately herein.

A "replicon" is any genetic element (e.g., plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*; i.e., capable of replication under its own

control.

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

5 A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in its either single stranded form, or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term  
10 includes double-stranded DNA found, *inter alia*, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes. In discussing the structure herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (i.e., the strand having a  
15 sequence homologous to the mRNA).

An "origin of replication" refers to those DNA sequences that participate in DNA synthesis.

A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory  
20 sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA,  
25 genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. A polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for the expression of a coding sequence in a host cell.

5 A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements  
10 necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site, as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.  
15 Eukaryotic promoters often, but not always, contain "TATA" boxes and "CAT" boxes. Prokaryotic promoters contain Shine-Dalgarno sequences in addition to the -10 and -35 consensus sequences.

An "expression control sequence" is a DNA sequence that controls and regulates the transcription and translation of another  
20 DNA sequence. A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then translated into the protein encoded by the coding sequence.

A "signal sequence" can be included near the coding  
25 sequence. This sequence encodes a signal peptide, N-terminal to the polypeptide, that communicates to the host cell to direct the polypeptide to the cell surface or secrete the polypeptide into the media, and this signal peptide is clipped off by the host cell

before the protein leaves the cell. Signal sequences can be found associated with a variety of proteins native to prokaryotes and eukaryotes.

The term "oligonucleotide", as used herein in referring to the probe of the present invention, is defined as a molecule comprised of two or more ribonucleotides, preferably more than three. Its exact size will depend upon many factors which, in turn, depend upon the ultimate function and use of the oligonucleotide.

The term "primer" as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product, which is complementary to a nucleic acid strand, is induced, i.e., in the presence of nucleotides and an inducing agent such as a DNA polymerase and at a suitable temperature and pH. The primer may be either single-stranded or double-stranded and must be sufficiently long to prime the synthesis of the desired extension product in the presence of the inducing agent. The exact length of the primer will depend upon many factors, including temperature, source of primer and use the method. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides.

The primers herein are selected to be "substantially" complementary to different strands of a particular target DNA sequence. This means that the primers must be sufficiently complementary to hybridize with their respective strands.

Therefore, the primer sequence need not reflect the exact sequence of the template. For example, a non-complementary nucleotide fragment may be attached to the 5' end of the primer, with the remainder of the primer sequence being complementary to the strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the primer, provided that the primer sequence has sufficient complementary with the sequence or hybridize therewith and thereby form the template for the synthesis of the extension product.

10 As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" by exogenous or heterologous DNA when such DNA has been introduced inside the cell.

15 The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the

20 transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of

25 cells derived from a single cell or ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

Two DNA sequences are "substantially homologous"

when at least about 75% (preferably at least about 80%, and most preferably at least about 90% or 95%) of the nucleotides match over the defined length of the DNA sequences. Sequences that are substantially homologous can be identified by comparing the sequences using standard software available in sequence data banks, or in a Southern hybridization experiment under, for example, stringent conditions as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Maniatis et al., *supra*; DNA Cloning, Vols. I & II, *supra*; Nucleic Acid Hybridization, *supra*.

A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another example, coding sequence is a construct where the coding sequence itself is not found in nature (e.g., a cDNA where the genomic coding sequence contains introns, or synthetic sequences having codons different than the native gene). Allelic variations or naturally-occurring mutational events do not give rise to a heterologous region of DNA as defined herein.

The labels most commonly employed for these studies are radioactive elements, enzymes, chemicals which fluoresce when exposed to ultraviolet light, and others. A number of fluorescent materials are known and can be utilized as labels. These include, for example, fluorescein, rhodamine, auramine, Texas Red, AMCA blue and Lucifer Yellow. A particular detecting material is anti-rabbit

antibody prepared in goats and conjugated with fluorescein through an isothiocyanate.

Proteins can also be labeled with a radioactive element or with an enzyme. The radioactive label can be detected by any of the currently available counting procedures. The preferred isotope may be selected from  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{36}\text{Cl}$ ,  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{58}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{90}\text{Y}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ , and  $^{186}\text{Re}$ .

Enzyme labels are likewise useful, and can be detected by any of the presently utilized colorimetric, spectrophotometric, fluorospectrophotometric, amperometric or gasometric techniques. The enzyme is conjugated to the selected particle by reaction with bridging molecules such as carbodiimides, diisocyanates, glutaraldehyde and the like. Many enzymes which can be used in these procedures are known and can be utilized. The preferred are peroxidase,  $\beta$ -glucuronidase,  $\beta$ -D-glucosidase,  $\beta$ -D-galactosidase, urease, glucose oxidase plus peroxidase and alkaline phosphatase. U.S. Patent Nos. 3,654,090, 3,850,752, and 4,016,043 are referred to by way of example for their disclosure of alternate labeling material and methods.

A particular assay system developed and utilized in the art is known as a receptor assay. In a receptor assay, the material to be assayed is appropriately labeled and then certain cellular test colonies are inoculated with a quantity of both the label after which binding studies are conducted to determine the extent to which the labeled material binds to the cell receptors. In this way, differences in affinity between materials can be ascertained.

An assay useful in the art is known as a "cis/trans" assay. Briefly, this assay employs two genetic constructs, one of which

is typically a plasmid that continually expresses a particular receptor of interest when transfected into an appropriate cell line, and the second of which is a plasmid that expresses a reporter such as luciferase, under the control of a receptor/ligand complex. Thus, for  
5 example, if it is desired to evaluate a compound as a ligand for a particular receptor, one of the plasmids would be a construct that results in expression of the receptor in the chosen cell line, while the second plasmid would possess a promoter linked to the luciferase gene in which the response element to the particular receptor is  
10 inserted. If the compound under test is an agonist for the receptor, the ligand will complex with the receptor, and the resulting complex will bind the response element and initiate transcription of the luciferase gene. The resulting chemiluminescence is then measured photometrically, and dose response curves are obtained and  
15 compared to those of known ligands. The foregoing protocol is described in detail in U.S. Patent No. 4,981,784.

As used herein, the term "host" is meant to include not only prokaryotes but also eukaryotes such as yeast, plant and animal cells. A recombinant DNA molecule or gene which encodes a human  
20 TADG-15 protein of the present invention can be used to transform a host using any of the techniques commonly known to those of ordinary skill in the art. Especially preferred is the use of a vector containing coding sequences for the gene which encodes a human TADG-15 protein of the present invention for purposes of prokaryote  
25 transformation. Prokaryotic hosts may include *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. Eukaryotic hosts include yeasts such as *Pichia pastoris*, mammalian cells and insect cells.



In general, expression vectors containing promoter sequences which facilitate the efficient transcription of the inserted DNA fragment are used in connection with the host. The expression vector typically contains an origin of replication, promoter(s), terminator(s), as well as specific genes which are capable of providing phenotypic selection in transformed cells. The transformed hosts can be fermented and cultured according to means known in the art to achieve optimal cell growth.

The invention includes a substantially pure DNA encoding a TADG-15 protein, a strand of which DNA will hybridize at high stringency to a probe containing a sequence of at least 15 consecutive nucleotides of (SEQ ID NO:1). The protein encoded by the DNA of this invention may share at least 80% sequence identity (preferably 85%, more preferably 90%, and most preferably 95%) with the amino acids listed in Figure 10 (SEQ ID NO:2). More preferably, the DNA includes the coding sequence of the nucleotides of Figure 9 (SEQ ID NO:1), or a degenerate variant of such a sequence.

The probe to which the DNA of the invention hybridizes preferably consists of a sequence of at least 20 consecutive nucleotides, more preferably 40 nucleotides, even more preferably 50 nucleotides, and most preferably 100 nucleotides or more (up to 100%) of the coding sequence of the nucleotides listed in Figure 9 (SEQ ID NO:1) or the complement thereof. Such a probe is useful for detecting expression of TADG-15 in a human cell by a method including the steps of (a) contacting mRNA obtained from the cell with the labeled hybridization probe; and (b) detecting hybridization of the probe with the mRNA.

This invention also includes a substantially pure

DNA containing a sequence of at least 15 consecutive nucleotides (preferably 20, more preferably 30, even more preferably 50, and most preferably all) of the region from nucleotides 1 to 3147 of the nucleotides listed in Figure 9 (SEQ ID NO:1).

5 By "high stringency" is meant DNA hybridization and wash conditions characterized by high temperature and low salt concentration, e.g., wash conditions of 65°C at a salt concentration of approximately 0.1 x SSC, or the functional equivalent thereof. For example, high stringency conditions may include hybridization at  
10 about 42°C in the presence of about 50% formamide; a first wash at about 65°C with about 2 x SSC containing 1% SDS; followed by a second wash at about 65°C with about 0.1 x SSC.

By "substantially pure DNA" is meant DNA that is not part of a milieu in which the DNA naturally occurs, by virtue of separation  
15 (partial or total purification) of some or all of the molecules of that milieu, or by virtue of alteration of sequences that flank the claimed DNA. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or  
20 eukaryote; or which exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by polymerase chain reaction (PCR) or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence, e.g., a fusion  
25 protein. Also included is a recombinant DNA which includes a portion of the nucleotides listed in Figure 9 (SEQ ID NO:1) which encodes an alternative splice variant of TADG-15.

The DNA may have at least about 70% sequence

identity to the coding sequence of the nucleotides listed in Figure 9 (SEQ ID NO:1), preferably at least 75% (e.g. at least 80%); and most preferably at least 90%. The identity between two sequences is a direct function of the number of matching or identical positions.

5 When a subunit position in both of the two sequences is occupied by the same monomeric subunit, e.g., if a given position is occupied by an adenine in each of two DNA molecules, then they are identical at that position. For example, if 7 positions in a sequence 10 nucleotides in length are identical to the corresponding positions

10 in a second 10-nucleotide sequence, then the two sequences have 70% sequence identity. The length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 100 nucleotides. Sequence identity is typically measured

15 using sequence analysis software (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705).

The present invention comprises a vector comprising a DNA sequence which encodes a human TADG-15 protein and said

20 vector is capable of replication in a host which comprises, in operable linkage: a) an origin of replication; b) a promoter; and c) a DNA sequence coding for said protein. Preferably, the vector of the present invention contains a portion of the DNA sequence shown in SEQ ID No:1. A "vector" may be defined as a replicable nucleic acid

25 construct, e.g., a plasmid or viral nucleic acid. Vectors may be used to amplify and/or express nucleic acid encoding TADG-15 protein. An expression vector is a replicable construct in which a nucleic acid sequence encoding a polypeptide is operably linked to suitable

control sequences capable of effecting expression of the polypeptide in a cell. The need for such control sequences will vary depending upon the cell selected and the transformation method chosen. Generally, control sequences include a transcriptional promoter and/or enhancer, suitable mRNA ribosomal binding sites, and sequences which control the termination of transcription and translation. Methods which are well known to those skilled in the art can be used to construct expression vectors containing appropriate transcriptional and translational control signals. See for example, the techniques described in Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual* (2nd Ed.), Cold Spring Harbor Press, N.Y. A gene and its transcription control sequences are defined as being "operably linked" if the transcription control sequences effectively control the transcription of the gene. Vectors of the invention include, but are not limited to, plasmid vectors and viral vectors. Preferred viral vectors of the invention are those derived from retroviruses, adenovirus, adeno-associated virus, SV40 virus, or herpes viruses.

By a "substantially pure protein" is meant a protein which has been separated from at least some of those components which naturally accompany it. Typically, the protein is substantially pure when it is at least 60%, by weight, free from the proteins and other naturally-occurring organic molecules with which it is naturally associated *in vivo*. Preferably, the purity of the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight. A substantially pure TADG-15 protein may be obtained, for example, by extraction from a natural source; by expression of a recombinant nucleic acid

encoding an TADG-15 polypeptide; or by chemically synthesizing the protein. Purity can be measured by any appropriate method, e.g., column chromatography such as immunoaffinity chromatography using an antibody specific for TADG-15, polyacrylamide gel electrophoresis, or HPLC analysis. A protein is substantially free of naturally associated components when it is separated from at least some of those contaminants which accompany it in its natural state. Thus, a protein which is chemically synthesized or produced in a cellular system different from the cell from which it naturally originates will be, by definition, substantially free from its naturally associated components. Accordingly, substantially pure proteins include eukaryotic proteins synthesized in *E. coli*, other prokaryotes, or any other organism in which they do not naturally occur.

In addition to substantially full-length proteins, the invention also includes fragments (e.g., antigenic fragments) of the TADG-15 protein (SEQ ID No:2). As used herein, "fragment," as applied to a polypeptide, will ordinarily be at least 10 residues, more typically at least 20 residues, and preferably at least 30 (e.g., 50) residues in length, but less than the entire, intact sequence. Fragments of the TADG-15 protein can be generated by methods known to those skilled in the art, e.g., by enzymatic digestion of naturally occurring or recombinant TADG-15 protein, by recombinant DNA techniques using an expression vector that encodes a defined fragment of TADG-15, or by chemical synthesis. The ability of a candidate fragment to exhibit a characteristic of TADG-15 (e.g., binding to an antibody specific for TADG-15) can be assessed by methods described herein. Purified TADG-15 or antigenic fragments of TADG-15 can be used to generate new antibodies or to

test existing antibodies (e.g., as positive controls in a diagnostic assay) by employing standard protocols known to those skilled in the art. Included in this invention are polyclonal antisera generated by using TADG-15 or a fragment of TADG-15 as the immunogen in, e.g., rabbits. Standard protocols for monoclonal and polyclonal antibody production known to those skilled in this art are employed. The monoclonal antibodies generated by this procedure can be screened for the ability to identify recombinant TADG-15 cDNA clones, and to distinguish them from known cDNA clones.

Further included in this invention are TADG-15 proteins which are encoded at least in part by portions of SEQ ID NO:2, e.g., products of alternative mRNA splicing or alternative protein processing events, or in which a section of TADG-15 sequence has been deleted. The fragment, or the intact TADG-15 polypeptide, may be covalently linked to another polypeptide, e.g. which acts as a label, a ligand or a means to increase antigenicity.

The invention also includes a polyclonal or monoclonal antibody which specifically binds to TADG-15. The invention encompasses not only an intact monoclonal antibody, but also an immunologically-active antibody fragment, e.g., a Fab or (Fab)<sub>2</sub> fragment; an engineered single chain Fv molecule; or a chimeric molecule, e.g., an antibody which contains the binding specificity of one antibody, e.g., of murine origin, and the remaining portions of another antibody, e.g., of human origin.

In one embodiment, the antibody, or a fragment thereof, may be linked to a toxin or to a detectable label, e.g. a radioactive label, non-radioactive isotopic label, fluorescent label, chemiluminescent label, paramagnetic label, enzyme

label, or colorimetric label. Examples of suitable toxins include diphtheria toxin, *Pseudomonas* exotoxin A, ricin, and cholera toxin. Examples of suitable enzyme labels include malate hydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, alcohol  
5 dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, acetylcholinesterase, etc. Examples of suitable radioisotopic labels  
10 include  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , etc.

Paramagnetic isotopes for purposes of *in vivo* diagnosis can also be used according to the methods of this invention. There are numerous examples of elements that are useful in magnetic resonance imaging. For discussions on *in vivo* nuclear magnetic  
15 resonance imaging, see, for example, Schaefer et al., (1989) *JACC* 14, 472-480; Shreve et al., (1986) *Magn. Reson. Med.* 3, 336-340; Wolf, G. L., (1984) *Physiol. Chem. Phys. Med. NMR* 16, 93-95; Wesbey et al., (1984) *Physiol. Chem. Phys. Med. NMR* 16, 145-155; Runge et al., (1984) *Invest. Radiol.* 19, 408-415. Examples of suitable fluorescent  
20 labels include a fluorescein label, an isothiocyalate label, a rhodamine label, a phycoerythrin label, a phycocyanin label, an allophycocyanin label, an ophthaldehyde label, a fluorescamine label, etc. Examples of chemiluminescent labels include a luminal label, an isoluminal label, an aromatic acridinium ester label, an imidazole  
25 label, an acridinium salt label, an oxalate ester label, a luciferin label, a luciferase label, an aequorin label, etc.

Those of ordinary skill in the art will know of other suitable labels which may be employed in accordance with

the present invention. The binding of these labels to antibodies or fragments thereof can be accomplished using standard techniques commonly known to those of ordinary skill in the art. Typical techniques are described by Kennedy et al., (1976) *Clin. Chim. Acta* 70, 1-31; and Schurs et al., (1977) *Clin. Chim. Acta* 81, 1-40. Coupling techniques mentioned in the latter are the glutaraldehyde method, the periodate method, the dimaleimide method, the m-maleimidobenzyl-N-hydroxy-succinimide ester method. All of these methods are incorporated by reference herein.

Also within the invention is a method of detecting TADG-15 protein in a biological sample, which includes the steps of contacting the sample with the labeled antibody, e.g., radioactively tagged antibody specific for TADG-15, and determining whether the antibody binds to a component of the sample.

As described herein, the invention provides a number of diagnostic advantages and uses. For example, the TADG-15 protein is useful in diagnosing cancer in different tissues since this protein is highly overexpressed in tumor cells. Antibodies (or antigen-binding fragments thereof) which bind to an epitope specific for TADG-15, are useful in a method of detecting TADG-15 protein in a biological sample for diagnosis of cancerous or neoplastic transformation. This method includes the steps of obtaining a biological sample (e.g., cells, blood, plasma, tissue, etc.) from a patient suspected of having cancer, contacting the sample with a labeled antibody (e.g., radioactively tagged antibody) specific for TADG-15, and detecting the TADG-15 protein using standard immunoassay techniques such as an ELISA. Antibody binding to the biological sample indicates that the sample contains a component which specifically binds to an epitope



within TADG-15.

Likewise, a standard Northern blot assay can be used to ascertain the relative amounts of TADG-15 mRNA in a cell or tissue obtained from a patient suspected of having cancer, in accordance with conventional Northern hybridization techniques known to those of ordinary skill in the art. This Northern assay uses a hybridization probe, e.g. radiolabelled TADG-15 cDNA, either containing the full-length, single stranded DNA having a sequence complementary to SEQ ID NO:1 (Figure 9), or a fragment of that DNA sequence at least 20 (preferably at least 30, more preferably at least 50, and most preferably at least 100 consecutive nucleotides in length). The DNA hybridization probe can be labeled by any of the many different methods known to those skilled in this art.

Antibodies to the TADG-15 protein can be used in an immunoassay to detect increased levels of TADG-15 protein expression in tissues suspected of neoplastic transformation. These same uses can be achieved with Northern blot assays and analyses.

The present invention is directed to DNA encoding a TADG-15 protein selected from the group consisting of: (a) isolated DNA which encodes a TADG-15 protein; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a TADG-15 protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to the degeneracy of the genetic code, and which encodes a TADG-15 protein. Preferably, the DNA has the sequence shown in SEQ ID No:1. More preferably, the DNA encodes a TADG-15 protein having the amino acid sequence shown in SEQ ID No:2.

The present invention is also directed to a

vector capable of expressing the DNA of the present invention adapted for expression in a recombinant cell and regulatory elements necessary for expression of the DNA in the cell. Preferably, the vector contains DNA encoding a TADG-15 protein having the amino acid  
5 sequence shown in SEQ ID No:2.

The present invention is also directed to a host cell transfected with the vector described herein, said vector expressing a TADG-15 protein. Representative host cells include consisting of bacterial cells, mammalian cells and insect cells.

10 The present invention is also directed to a isolated and purified TADG-15 protein coded for by DNA selected from the group consisting of: (a) isolated DNA which encodes a TADG-15 protein; (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a TADG-15 protein; and (c) isolated DNA differing from the  
15 isolated DNAs of (a) and (b) above in codon sequence due to the degeneracy of the genetic code, and which encodes a TADG-15 protein. Preferably, the isolated and purified TADG-15 protein of claim 9 having the amino acid sequence shown in SEQ ID No:2.

The present invention is also directed to a method of  
20 detecting expression of the protein of claim 1, comprising the steps of: (a) contacting mRNA obtained from the cell with the labeled hybridization probe; and (b) detecting hybridization of the probe with the mRNA.

The following examples are given for the purpose of  
25 illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

### EXAMPLE 1

#### 5 Tissue collection and storage

Upon patient hysterectomy, bilateral salpingo-oophorectomy, or surgical removal of neoplastic tissue, the specimen is retrieved and placed it on ice. The specimen was then taken to the resident pathologist for isolation and identification of specific tissue  
10 samples. Finally, the sample was frozen in liquid nitrogen, logged into the laboratory record and stored at -80°C. Additional specimens were frequently obtained from the Cooperative Human Tissue Network (CHTN). These samples were prepared by the CHTN and shipped to us on dry ice. Upon arrival, these specimens were logged into the  
15 laboratory record and stored at -80°C.

### EXAMPLE 2

#### 20 mRNA isolation and cDNA synthesis

Forty-one ovarian tumors (10 low malignant potential tumors and 31 carcinomas) and 10 normal ovaries were obtained from surgical specimens and frozen in liquid nitrogen. The human ovarian carcinoma cell lines SW 626 and Caov 3, the human breast  
25 carcinoma cell lines MDA-MB-231 and MDA-MB-435S, and the human uterine cervical carcinoma cell line Hela were purchased from the American Type Culture Collection (Rockville, MD). Cells were cultured to subconfluency in Dulbecco's modified Eagle's medium,

suspended with 10% (v/v) fetal bovine serum and antibiotics.

Messenger RNA (mRNA) isolation was performed according to the manufacturer's instructions using the Mini RiboSep™ Ultra mRNA isolation kit purchased from Becton Dickinson (cat. # 30034). This was an oligo(dt) chromatography based system of mRNA isolation. The amount of mRNA recovered was quantitated by UV spectrophotometry.

First strand complementary DNA (cDNA) was synthesized using 5.0 mg of mRNA and either random hexamer or oligo(dT) primers according to the manufacturer's protocol utilizing a first strand synthesis kit obtained from Clontech (cat.# K1402-1). The purity of the cDNA was evaluated by PCR using primers specific for the p53 gene. These primers span an intron such that pure cDNA can be distinguished from cDNA that is contaminated with genomic DNA.

### EXAMPLE 3

#### PCR reactions

The mRNA overexpression of TADG-15 was determined using a quantitative PCR. Oligonucleotide primers were used for: TADG-15, forward 5'-ATGACAGAGGATTCAGGTAC-3' (SEQ ID NO: 10) and reverse 5'-GAAGGTGAAGTCATTGAAGA-3' (SEQ ID NO: 11); and  $\beta$ -tubulin, forward 5'-TGCATTGACAACGAGGC-3' (SEQ ID NO: 12) and reverse 5'-CTGTCTTGACATTGTTG-3' (SEQ ID NO: 13).  $\beta$ -tubulin was utilized as an internal control. Reactions were carried out as follows: first strand cDNA generated from 50 ng of mRNA will be used as template in the presence of 1.0 mM MgCl<sub>2</sub>, 0.2 mM dNTPs, 0.025 U Taq polymerase/ml of reaction, and 1 x buffer supplied with

enzyme. In addition, primers must be added to the PCR reaction. Degenerate primers which may amplify a variety of cDNAs are used at a final concentration of 2.0 mM each, whereas primers which amplify specific cDNAs are added to a final concentration of 0.2 mM each.

After initial denaturation at 95°C for 3 minutes, thirty cycles of PCR are carried out in a Perkin Elmer Gene Amp 2400 thermal cycler. Each cycle consists of 30 seconds of denaturation at 95°C, 30 seconds of primer annealing at the appropriate annealing temperature, and 30 seconds of extension at 72°C. The final cycle will be extended at 72°C for 7 minutes. To ensure that the reaction succeeded, a fraction of the mixture will be electrophoresed through a 2% agarose/TAE gel stained with ethidium bromide (final concentration 1 mg/ml). The annealing temperature varies according to the primers that are used in the PCR reaction. For the reactions involving degenerate primers, an annealing temperature of 48°C were used. The appropriate annealing temperature for the TADG-15 and  $\beta$ -tubulin specific primers is 62°C.

#### EXAMPLE 4

##### T-vector ligation and transformations

The purified PCR products are ligated into the Promega T-vector plasmid and the ligation products are used to transform JM109 competent cells according to the manufacturer's instructions (Promega cat. #A3610). Positive colonies were cultured for amplification, the plasmid DNA isolated by means of the Wizard<sup>TM</sup> Minipreps DNA purification system (Promega cat #A7500), and the plasmids were

digested with ApaI and SacI restriction enzymes to determine the size of the insert. Plasmids with inserts of the size(s) visualized by the previously described PCR product gel electrophoresis were sequenced.

5

### EXAMPLE 5

#### DNA sequencing

Utilizing a plasmid specific primer near the cloning site, sequencing reactions were carried out using PRISM<sup>TM</sup> Ready Reaction Dye Deoxy<sup>TM</sup> terminators (Applied Biosystems cat# 401384) according to the manufacturer's instructions. Residual dye terminators were removed from the completed sequencing reaction using a Centri-sep<sup>TM</sup> spin column (Princeton Separation cat.# CS-901). An Applied Biosystems Model 373A DNA Sequencing System was available and was used for sequence analysis. Based upon the determined sequence, primers that specifically amplify the gene of interest were designed and synthesized.

15

### EXAMPLE 6

20

#### Northern blot analysis

10 µg mRNAs were size separated by electrophoresis through a 1% formaldehyde-agarose gel in 0.02 M MOPS, 0.05 M sodium acetate (pH 7.0), and 0.001 M EDTA. The mRNAs were then blotted to Hybond-N (Amersham) by capillary action in 20 x SSPE. The RNAs are fixed to the membrane by baking for 2 hours at 80°C. Additional multiple tissue northern (MTN) blots were purchased from CLONTECH Laboratories, Inc. These blots include the Human

25

MTN blot (cat.#7760-1), the Human MTN II blot (cat.#7759-1), the Human Fetal MTN II blot (cat.#7756-1), and the Human Brain MTN III blot (cat.#7750-1). The appropriate probes were radiolabelled utilizing the Prime-a-Gene Labeling System available from Promega (cat#U1100). The blots were probed and stripped according to the ExpressHyb Hybridization Solution protocol available from CLONTECH (cat.#8015-1 or 8015-2).

#### EXAMPLE 7

##### 10 Quantitative PCR

Quantitative-PCR was performed in a reaction mixture consisting of cDNA derived from 50 ng of mRNA, 5 pmol of sense and antisense primers for TADG-15 and the internal control  $\beta$ -tubulin, 0.2 mmol of dNTPs, 0.5 mCi of [ $\alpha$ - $^{32}$ P]dCTP, and 0.625 U of Taq polymerase in 1 x buffer in a final volume of 25  $\mu$ l. This mixture was subjected to 1 minute of denaturation at 95°C followed by 30 cycles of denaturation for 30 seconds at 95°C, 30 seconds of annealing at 62°C, and 1 minute of extension at 72°C with an additional 7 minutes of extension on the last cycle. The product was electrophoresed through a 2% agarose gel for separation, the gel was dried under vacuum and autoradiographed. The relative radioactivity of each band was determined by PhosphorImager from Molecular Dynamics.

25

#### EXAMPLE 8

The present invention describes the use of primers directed to conserved areas of the serine protease family to

identify members of that family which are overexpressed in carcinoma. Several genes were identified and cloned in other tissues, but not previously associated with ovarian carcinoma. The present invention describes a protease identified in ovarian carcinoma. This  
5 gene was identified using primers to the conserved area surrounding the catalytic domain of the conserved amino acid histidine and the downstream conserved amino acid serine which lies approximately 150 amino acids towards the carboxyl end of the protease.

The gene encoding the novel extracellular serine protease  
10 of the present invention was identified from a group of proteases overexpressed in carcinoma by subcloning and sequencing the appropriate PCR products. An example of such a PCR reaction is given in Figure 1. Subcloning and sequencing of individual bands from such an amplification provided a basis for identifying the protease of the  
15 present invention.

### EXAMPLE 9

The sequence determined for the catalytic domain of  
20 TADG-15 is presented in Figure 2 and is consistent with other serine proteases and specifically contains conserved amino acids appropriate for the catalytic domain of the trypsin-like serine protease family. Specific primers (20mers) derived from this sequence were used.

A series of normal and tumor cDNAs were examined to  
25 determine the expression of the TADG-15 gene in ovarian carcinoma. In a series of normal derived cDNA compared to carcinoma derived cDNA using  $\beta$ -tubulin as an internal control for PCR amplification, TADG-15 was significantly overexpressed in all of the



carcinomas examined and either was not detected or was detected at a very low level in normal epithelial tissue (Figure 3). This evaluation was extended to a standard panel of about 40 tumors. Using these specific primers, the expression of this gene was also examined in  
5 tumor cell lines derived from both ovarian and breast carcinoma tissues as shown in Figure 5 and in other tumor tissues as shown in Figure 6. The expression of TADG-15 was also observed in carcinomas of the breast, colon, prostate and lung.

Using the specific sequence for TADG-15 covering the full  
10 domain of the catalytic site as a probe for Northern blot analysis, three Northern blots were examined: one derived from ovarian tissues, both normal and carcinoma; one from fetal tissues; and one from adult normal tissues. As shown in Figure 7, TADG-15 transcripts were noted in all ovarian carcinomas, but were not present in  
15 detectable levels in any of the following tissues: a) normal ovary, b) fetal liver and brain, c) adult spleen, thymus, testes, ovary and peripheral blood lymphocytes, d) skeletal muscle, liver, brain or heart. The transcript size was found to be approximately 3.2 kb. The hybridization for the fetal and adult blots was appropriate and done  
20 with the same probe as with the ovarian tissue. Subsequent to this examination, it was confirmed that these blots contained other detectable mRNA transcripts

Initially using the catalytic domain of the protease to probe Hela cDNA and ovarian tumor cDNA libraries, one clone was  
25 obtained covering the entire 3' end of the TADG-15 gene from the ovarian tumor library. On further screening using the 5' end of the newly detected clones, two more clones were identified covering the 5' end of the TADG-15 gene from the Hela library (Figure 8). The

complete nucleotide sequence (SEQ ID No:1) is provided in Figure 9 along with translation of the open reading frame (SEQ ID No:2).

In the nucleotide sequence, there is a Kozak sequence typical of sequences upstream from the initiation site of translation.

5 There is also a poly-adenylation signal sequence and a poly-adenylated tail. The open reading frame consists of a 855 amino acid sequence (SEQ ID No:2) which includes an amino terminal cytoplasmic tail from amino acids 1-50, an approximately 22 amino acid transmembrane domain followed by an extracellular sequence

10 preceding two CUB repeats identified from complement subcomponents C1r and C1s. These two repeats are followed by four repeat domains of a class A motif of the LDL receptor and these four repeats are followed by the protease enzyme of the trypsin family constituting the carboxyl end of the TADG-15 protein (Figure 11).

15 Also a clear delineation of the catalytic domain conserved histidine, aspartic acid, serine series along with a series of amino acids conserved in the serine protease family is indicated (Figure 10).

A search of GeneBank for similar previously identified sequences yielded one such sequence with relatively high homology

20 to a portion of the TADG-15 gene. The similarity between the portion of TADG-15 from nucleotide #182 to 3139 and SNC-19 (SEQ ID No: 9; GeneBank accession #U20428) is approximately 97% (Figure 12). There are however significant differences between SNC-19 and TADG-15 viz. TADG-15 has an open reading frame of 855 amino acids

25 whereas the longest ORF of SNC-19 is only 173 amino acids. SNC-19 does not include a proper start site for the initiation of translation nor does it include the amino terminal portion of the protein encoded by TADG-15. Moreover, SNC-19 does not include an ORF for a

functional serine protease because the His, Asp and Ser residues necessary for function are encoded in different reading frames.

TADG-15 is a highly overexpressed gene in tumors. It is expressed in a limited number of normal tissues, primarily tissues that are involved in either uptake or secretion of molecules e.g. colon and pancreas. TADG-15 is further novel in its component structure of domains in that it has a protease catalytic domain which could be released and used as a diagnostic and which has the potential for a target for therapeutic intervention. TADG-15 also has ligand binding domains which are commonly associated with molecules that internalize or take-up ligands from the external surface of the cell as does the LDL receptor for the LDL cholesterol complex. There is potential that these domains may be involved in uptake of specific ligands and they may offer the potential for making delivery of toxic molecules or genes to tumor cells which express this molecule on their surface. It has features that are similar to the hepsin serine protease molecule in that it also has an amino-terminal transmembrane domain with the proteolytic catalytic domain extended into the extracellular matrix. The difference here is that TADG-15 includes these ligand binding repeat domains which the hepsin gene does not have. In addition to the use of this gene as a diagnostic or therapeutic target in ovarian carcinoma and other carcinomas including breast, prostate, lung and colon, its ligand-binding domains may be valuable in the uptake of specific molecules into tumor cells.

Any patents or publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. These patents and publications are herein incorporated by reference to the same extent as

if each individual publication was specifically and individually indicated to be incorporated by reference.

One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects and obtain  
5 the ends and advantages mentioned, as well as those inherent therein. The present examples along with the methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention.  
10 Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

## WHAT IS CLAIMED IS:

1. DNA encoding a TADG-15 protein selected from the  
5 group consisting of:
  - (a) isolated DNA which encodes a TADG-15 protein;
  - (b) isolated DNA which hybridizes to isolated DNA of (a)  
above and which encodes a TADG-15 protein; and
  - (c) isolated DNA differing from the isolated DNAs of (a)  
10 and (b) above in codon sequence due to the degeneracy of the genetic  
code, and which encodes a TADG-15 protein.
2. The DNA of claim 1, wherein said DNA has the  
15 sequence shown in SEQ ID No:1.
3. The DNA of claim 1, wherein said TADG-15 protein  
has the amino acid sequence shown in SEQ ID No:2.  
20
4. A vector capable of expressing the DNA of claim  
1 adapted for expression in a recombinant cell and regulatory  
elements necessary for expression of the DNA in the cell.  
25
5. The vector of claim 4, wherein said DNA encodes a  
TADG-15 protein having the amino acid sequence shown in SEQ ID  
No:2.

6. A host cell transfected with the vector of claim 4, said vector expressing a TADG-15 protein.

5

7. The host cell of claim 6, wherein said cell is selected from group consisting of bacterial cells, mammalian cells, plant cells and insect cells.

10

8. The host cell of claim 7, wherein said bacterial cell is *E. coli*.

15

9. Isolated and purified TADG-15 protein coded for by DNA selected from the group consisting of:

- (a) isolated DNA which encodes a TADG-15 protein;
- (b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a TADG-15 protein; and
- (c) isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to the degeneracy of the genetic code, and which encodes a TADG-15 protein.

20

10. The isolated and purified TADG-15 protein of claim 9 having the amino acid sequence shown in SEQ ID No:2.

11. A method of detecting expression of the protein of claim 1, comprising the steps of:

(a) contacting mRNA obtained from the cell with the labeled hybridization probe; and

5 (b) detecting hybridization of the probe with the mRNA.




FIG. 1



RIVGGRDTSI GRWPQVSL. ....RYDG.A HLCGSLLSG DWILTAACHF PE....RNRV LSRWRVFAGA VAQASPHGLC  
 RVVGGTDADE GEWPQVSL. ....HALQG HICGASLISP NWLVSAACHY IDDRGERYS D PTQWTAFLGL HDQSORSAPC  
 KIIDGAPCAR GSHPWQVAL. ....LSGNQL H.CGGVLVNE RWILTAACH. ....K MNEYTVHLGS DTLG...DR.F  
 KIVGGYNCEE NSVPYQVSL. ....NSGYHF .CGGSLINE QWVVSAGHC. ....Y KSRIQVRLGE HNIEVLEG.F  
 RIVNGEDAVP GSWPWQVSL. ....QDKTGF HFCGGLISE DWVVTAAHC. ....GV RTSDDVVAGE FDQGSDEE.F  
 RIVGGKVCCK GECPWQVLL. ....LVNG.A QLCGGTLINT IWVVSAAHCF DKIKNWRNLI .....AVLGE HDLSEHDGDI  
 RIKGGLFADI ASHPWQAAIF AKHRRSPGER FLCGGILISS CWILSAACHF QERFPPHLL. ....TVILGR .TYRUVVPEI

LGVQAVVYHG GYLPRDPNS EENSNDIALV HLSS.PLPLT EYIQPVCLPA ...AGQALVD GKICTVTGWS NTQYYGQQ.1  
 VQERRLKRII SHPFENDTFE D...YDIALI ELEK.PAEYS SMVRPICLPD ...ASHVFA GKAIWVTGWS HTQYGGTG.1  
 AQRIKASKSF RHPGYSTQT. ..HVNDMLV KLNS.QARLS SMVKVRLPS ...RCE..PP GTTCTVSGWG TTTSPDVTFI  
 EQFINAAKII RHPQYDRKT. ..LNNDIMLI KLSS.RAVIN ARVSTISLPT ...APP..AT GTKCLISGWG NTASSGADYI  
 IQVLKIAKVF KNPKEFILT. ..VNNDITLL KLAT.PARES QTVSAVCLPS ...ADDDEPA GTLCATTGWS KTKYNANKTI  
 QSRRAQVVI P....STYVP GTTNHDIALL RLHQ.PVULT DHVVPICLPE RTFESERTLAF VRFSLVSGWG QLIDRGATAI  
 EQKFEVEKYI VHKEFDDDTY D...NDIALI QLKSDSSRCA QESSVVRTVC LPPADLQLPD WTECELSGYG KHEALSPEFY:

GVLQEARVPI ISNDVCNGAD FYGN..QIKP KMFCAGYPEG G.....IDA CQGDSSGGPFV CEDSISRTPR WRLCGIVSWI  
 LILQKGEIRV INQTTCE..N LIPQ..QITP RMMCVGFSLG G.....VDS CQGDSSGGPL. ..SSVEADGR IFQAGVVSWI  
 SLMCVDVKL ISPODCTKV. .YKD..LLEN SMLCAGIPDS K.....KNA CNGDSSGGPLV C....R.... GTLQGLVSWI  
 DELOCLDAPV LSQAKCEAS. .YPG..KITS NMFCVGFLEG G.....KDS CQGDSSGGPVV C....N.... GQLQGVVSWI  
 DKLQQAALPL LSNAECKKS. .WGR..RITD VMICAG..AS G.....VSS CMGDSSGGPLV C....QKOGA WTLVGIVSWI  
 ELMVLNVPRIL MTQDCLQQR KVGDSNPITE YMFCAGYS DG S.....KDS CKGDSSGGP.. ..HATHYRGT WYLTGIVSWI  
 ERLKEAHVRL YPSSRCTSQH LLNRT..VTD NMLCAGDTRS GGQANLHDA CQGDSSGGPLV CLN....DGR MTLVGIISWI

T.GCALAQKP GYTKVSDER EWIFQAIKTH SEASGMVTQL ~ (SEQ. ID NO: 3) Heps  
 D.GCAQRNKP GYTRLPLFR DWIKENTGV~ ~ (SEQ. ID NO: 14) Tadg 15  
 TFPCGQPNDP GYTVQVCKFT KWINDTMKKH R~ ~ (SEQ. ID NO: 4) Scce  
 D.GCAQKNKP GYTKVYNYV KWIKNTIAAN S~ ~ (SEQ. ID NO: 5) Try  
 SDTCS.TSSP GVIARVTKLI PWVQKILAAAN ~ ~ (SEQ. ID NO: 6) Chymb  
 Q.GCATVGHE GYTRVVSQYI EWLQKLMRSE PRPGVLLRAP FP (SEQ. ID NO: 7) Fac 7  
 .LGGQKQKDV GYTKVVTNYL DWIRDNMRP~ ~ (SEQ. ID NO: 8) Tpa

FIG. 2




FIG. 3




Figure 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8





1 MGSDRARKGG GGPKDFGAGL KYNSRHEKVN GLEEGVEFLP VNNVKKVEKH 1  
 51 GPGRVWVLAA VLIGLLLVLL GIGFLVWHLQ YRDVRVQKVF NGYMRITNEN 2  
 101 FVDAYENS NS TEFVSLASKV KDALKLLYSG VPFLGPHYKE SAVTAFSEGS  
 151 VIAYYWSEFS IPQHLVEEAE RVMAEERVVM LPPRARSLSK FVVTSVVAFR  
 201 TDSKTVQRTQ DNS<sup>\*</sup>CSFGLHA RGVELMRFTT PGFPDSPYPA HAR<sup>\*</sup>CQWALRG  
 251 DADSVLSLTF RSFDLAS<sup>\*</sup>CDE RGS<sup>\*</sup>DLVTYVY TLSPMEPHAL VQL<sup>\*</sup>CGTYPPS  
 301 YNLT<sup>\*</sup>FHSSQN VLLITLITNT ERRHPGFEAT FFQLPRMSS<sup>\*</sup> GGRLRKAQGT 3  
 351 FNSPYYPGHY PPNID<sup>\*</sup>CTWNI EVENNQHVKV SFKFFYLLEP GVPAGT<sup>\*</sup>CPKD  
 401 YVEINGEKYC<sup>\*</sup> GERSQFVVTS NSNKITVRFH SDQSYTDTGF LAEYLSY<sup>\*</sup>DSS  
 451 DPCPGQFTCR TGR CIRKELR CDGWADCTDH SDE<sup>\*</sup>LNCSCDA GHQFTCKNKF  
 501 CKPLFWVCD S VND CGDN<sup>\*</sup>SDE QGCSCPAQTF RCSNGKCLSK SQQCNGKDDC 4  
 551 GDG<sup>\*</sup>SDEASCP KVNVTCTKH TYRCLNGLCL SKGNPECDGK EDCSDG<sup>\*</sup>SDEK  
 601 DCD<sup>\*</sup>CGLRST RQARVVGTD ADEGEWPWQV SLHALGQGH I CGASLISP NW  
 651 LVSAAH<sup>\*</sup>CYID DRGFRYS DPT QWTAFLGLHD QSQRSAPGVQ ERRLKRIISH  
 701 PFFNDFTFDY <sup>○</sup>IALLELEKP AEYSSMVRPI CLPDASHVFP AGKAIWVTGW 5  
 751 GHTQYGGTGA LILQKGEIRV INQTT CENLL PQQITPRMMC VGFLSGGVDS  
 801 CQGL<sup>\*</sup>SGGPLS SVEADGRIFQ AGVVS<sup>\*</sup>WGDGC AQRNKP<sup>\*</sup>GVYT RLPLFRDWIK  
 851 ENTGV (SEQ. ID NO: 2)

\* : Conserved cysteine residue

NXT : Possible N-linked glycosylation site

SDE : Conserved SDE motif


○ : Potential cleavage site

: Conserved amino acids of catalytic triad H, D, S

1. Cytoplasmic domain
2. Transmembrane domain
3. CUB repeat
4. Ligand-binding repeat (class A motif)  
of LDL receptor like domain
5. Serine protease

FIG. 10

11/13



1. Cytoplasmic domain
2. Transmembrane domain
3. Extracellular domain
- 4-5. CUB repeat
- 6-9. Ligand-binding repeat (class A motif) of LDL receptor like domain
10. Serine protease

FIG. 11

12/13

```

LOCUS       HSU20428            2900 bp      mRNA
DEFINITION  Human SNC19 mRNA sequence.
ACCESSION   U20428
NID         g1890631
KEYWORDS
SOURCE      human.
ORGANISM    Homo sapiens
             Eukaryota; Eukaryotes; Metazoa; Chordata;
             Vertebrata; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE   1 (bases 1 to 2900)
AUTHORS     Zheng, S., Cai, X., Gang, L., Cao, J., Zheng, L. and Zhi, Z.J.
TITLE       SNC19 gene in Homo sapiens
JOURNAL     Unpublished
REFERENCE   2 (bases 1 to 2900)
AUTHORS     Zheng, S.
TITLE       Direct Submission
JOURNAL     Submitted (20-JAN-1995) Shu Zheng, Jinder Institute, Zhejiang
             Medical University, Hangzhou, 310003, Peoples Republic of China

```

TADG15: TCACAGCGGGCCTCGGGGTACCATCGGG:10217CGTCCCTGGACCTGGGAGGGGGCCCGAAGCACTTCGGGGCCGGACT 81

SEC19: .....

[illegible]

Figure 12



## SEQUENCE LISTING

<110> O'Brien, Timothy J.  
 Tanimoto, Hirotoshi  
 <120> TADG-15: An Extracellular Serine Protease  
 Overexpressed in Breast and Ovarian Carcinomas  
 <130> D6064PCT  
 <140> PCT/US99/03436  
 <141> 1999-02-18  
 <150> US 09/027,337  
 <151> 1998-02-20  
 <160> 14  
 <210> 1  
 <211> 3147  
 <212> DNA  
 <213> *Homo sapiens*  
 <220>  
 <223> cDNA sequence of TADG-15  
 <400> 1

```

tcaagagcgg cctcggggta ccatggggag cgatcggggc cgcaagggcg 50
gagggggccc gaaggacttc ggcgcgggac tcaagtacaa ctcccggcac 100
gagaaagtga atggcttgga ggaaggcgtg gatttctgc cagtcaacaa 150
cgtcaagaag gtggaaaagc atggcccggg gcgctgggtg gtgctggcag 200
ccgtgctgat cggcctcctc ttggtcttgc tggggatcgg cttcctgggtg 250
tggcatttgc agtaccggga cgtgcgtgtc cagaaggtct tcaatggcta 300
catgaggatc acaaattgaga attttgtgga tgcctacgag aactccaact 350
ccactgagtt tgtaagcctg gccagcaagg tgaaggacgc gctgaagctg 400
ctgtacagcg gagtcccatt cctggggccc taccacaagg agtcggctgt 450
gacggccttc agcgagggca gcgtcatcgc ctactactgg tctgagttca 500
gcatcccgca gcacctggtg gaggaggcgg agcgcgtcat ggccgaggag 550
cgcgtagtca tgetgcccc gggggcgcg cccctgaagt cctttgtggt 600
cacctcagtg gtggctttcc ccacggactc caaaacagta cagaggaccc 650
aggacaacag ctgcagcttt ggctgcacg cccgcggtgt ggagctgatg 700
cgcttcacca cgcccggctt ccctgacagc cctaccccg ctcattgccc 750
ctgccagtg gcccgtggg gggacgcca ctcagtgtg agcctcacct 800
tccgcagctt tgaccttgcg tctgacgag agcgcggcag cgacctgggtg 850
acggtgtaca acacctgag ccccatggag cccacgccc tgggtgcagtt 900
gtgtggcacc taccctccct cctacaacct gacctccac tctcccaga 950
acgtcctgct catcacactg ataaccaaca ctgagcggcg gcatcccggc 1000
  
```

tttgaggcca ccttcttcca gctgcctagg atgagcagct gtggaggccg 1050  
cttacgtaaa gcccagggga cattcaacag cccctactac ccaggccact 1100  
acccacccaa cattgactgc acatggaaca ttgaggtgcc caacaaccag 1150  
catgtgaagg tgagcttcaa attcttctac ctgctggagc cggcggtgcc 1200  
tgcgggcacc tgccccaagg actacgtgga gatcaatggg gagaaatact 1250  
gcgagagagag gtcccagttc gtcgtcacca gcaacagcaa caagatcaca 1300  
gttcgcttcc actcagatca gtccctacacc gacaccggct tcttagctga 1350  
atacctctcc tacgactcca gtgacccatg cccggggcag ttcacgtgcc 1400  
gcacggggcg gtgtatccgg aaggagctgc gctgtgatgg ctgggcccag 1450  
tgcaccgacc acagcgatga gctcaactgc agttgcgacg ccggccacca 1500  
gttcacgtgc aagaacaagt tctgcaagcc cctcttctgg gtctgcgaca 1550  
gtgtgaacga ctgcggagac aacagcgacg agcaggggtg cagttgtccg 1600  
gcccagacct tcaggtgttc caatgggaag tgcctctcga aaagccagca 1650  
gtgcaatggg aaggacgact gtggggacgg gtccgacgag gcctcctgcc 1700  
ccaaggtgaa cgtcgtcact tgtaccaaac acacctaccg ctgcctcaat 1750  
gggctctgct tgagcaaggg caaccctgag tgtgacggga aggaggactg 1800  
tagcgacggc tcagatgaga aggactgcca ctgtgggctg cggtcattca 1850  
cgagacaggc tcgtgttgtt gggggcacgg atgcggatga gggcgagtgg 1900  
ccctggcagg taagcctgca tgctctgggc cagggccaca tctgcggtgc 1950  
ttccctcacc tctcccaact ggctggtctc tgccgcacac tgctacatcg 2000  
atgacagagg attcaggtac tcagacccca cgcagtggac ggccctcctg 2050  
ggcttgacag accagagcca gcgcagcgcc cctgggggtgc aggagcgcag 2100  
gctcaagcgc atcatctccc accccttctt caatgacttc accttcgact 2150  
atgacatcgc gctgctggag ctggagaaac cggcagagta cagctccatg 2200  
gtgcggccca tctgcctgcc ggacgcctcc catgtcttcc ctgccggcaa 2250  
ggccatcttg gtccacgggt ggggacacac ccagtatgga ggcactggcg 2300  
cgctgatcct gcaaaagggg gagatccgcg tcatcaacca gaccacctgc 2350  
gagaacctcc tgccgcagca gatcacgccg cgcattgatgt gcgtgggctt 2400  
cctcagcggc ggcggtggact cctgccaggg tgattccggg ggaccctgt 2450  
ccagcgtgga ggcggtatgg cggatcttcc aggcgggtgt ggtgagctgg 2500  
ggagacggct gcgctcagag gaacaagcca ggctgtgaca caaggctccc 2550  
tctgtttcgg gactggatca aagagaacac tggggtatag gggccggggc 2600  
caccctaatg tgtacacctg cggggccacc catcgtccac ccagtggtgc 2650  
acgcctgcag gctggagact ggaccgctga ctgcaccagc gccccagaa 2700  
catacactgt gaactcaatc tccagggtc caaatctgcc tagaaaacct 2750  
ctcgtctcct cagcctccaa agtggagctg ggaggtagaa ggggaggaca 2800  
ctgggtggttc tactgacca actgggggca aaggtttgaa gacacagcct 2850  
cccccgccag cccaagctg ggccgaggcg cgtttgtgta tatctgcctc 2900

ccctgtctgt aaggagcagc gggaaacggag cttcggagcc tcttcagtga 2950  
 aggtgggtggg gctgccggat ctgggctgtg gggcccttgg gccacgctct 3000  
 tgaggaagcc caggctcgga ggaccctgga aaacagacgg gtctgagact 3050  
 gaaattgttt taccagctcc caggggtggac ttcagtgtgt gtatttgtgt 3100  
 aaatgggtaa aacaatttat ttcttttttaa aaaaaaaaaa aaaaaaa 3147

<210> 2  
 <211> 855  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <223> Amino acid sequence of TADG-15 encoded by cDNA.  
 <400> 2

Met Gly Ser Asp Arg Ala Arg Lys Gly Gly Gly Gly Pro Lys Asp		
	5	10 15
Phe Gly Ala Gly Leu Lys Tyr Asn Ser Arg His Glu Lys Val Asn		
	20	25 30
Gly Leu Glu Glu Gly Val Glu Phe Leu Pro Val Asn Asn Val Lys		
	35	40 45
Lys Val Glu Lys His Gly Pro Gly Arg Trp Val Val Leu Ala Ala		
	50	55 60
Val Leu Ile Gly Leu Leu Leu Val Leu Leu Gly Ile Gly Phe Leu		
	65	70 75
Val Trp His Leu Gln Tyr Arg Asp Val Arg Val Gln Lys Val Phe		
	80	85 90
Asn Gly Tyr Met Arg Ile Thr Asn Glu Asn Phe Val Asp Ala Tyr		
	95	100 105
Glu Asn Ser Asn Ser Thr Glu Phe Val Ser Leu Ala Ser Lys Val		
	110	115 120
Lys Asp Ala Leu Lys Leu Leu Tyr Ser Gly Val Pro Phe Leu Gly		
	125	130 135
Pro Tyr His Lys Glu Ser Ala Val Thr Ala Phe Ser Glu Gly Ser		
	140	145 150
Val Ile Ala Tyr Tyr Trp Ser Glu Phe Ser Ile Pro Gln His Leu		
	155	160 165
Val Glu Glu Ala Glu Arg Val Met Ala Glu Glu Arg Val Val Met		
	170	175 180

Leu Pro Pro Arg	Ala Arg Ser Leu Lys Ser Phe Val Val Thr Ser	185	190	195
Val Val Ala Phe	Pro Thr Asp Ser Lys Thr Val Gln Arg Thr Gln	200	205	210
Asp Asn Ser Cys	Ser Phe Gly Leu His Ala Arg Gly Val Glu Leu	215	220	225
Met Arg Phe Thr	Thr Pro Gly Phe Pro Asp Ser Pro Tyr Pro Ala	230	235	240
His Ala Arg Cys	Gln Trp Ala Leu Arg Gly Asp Ala Asp Ser Val	245	250	255
Leu Ser Leu Thr	Phe Arg Ser Phe Asp Leu Ala Ser Cys Asp Glu	260	265	270
Arg Gly Ser Asp	Leu Val Thr Val Tyr Asn Thr Leu Ser Pro Met	275	280	285
Glu Pro His Ala	Leu Val Gln Leu Cys Gly Thr Tyr Pro Pro Ser	290	295	300
Tyr Asn Leu Thr	Phe His Ser Ser Gln Asn Val Leu Leu Ile Thr	305	310	315
Leu Ile Thr Asn	Thr Glu Arg Arg His Pro Gly Phe Glu Ala Thr	320	325	330
Phe Phe Gln Leu	Pro Arg Met Ser Ser Cys Gly Gly Arg Leu Arg	335	340	345
Lys Ala Gln Gly	Thr Phe Asn Ser Pro Tyr Tyr Pro Gly His Tyr	350	355	360
Pro Pro Asn Ile	Asp Cys Thr Trp Asn Ile Glu Val Pro Asn Asn	365	370	375
Gln His Val Lys	Val Ser Phe Lys Phe Phe Tyr Leu Leu Glu Pro	380	385	390
Gly Val Pro Ala	Gly Thr Cys Pro Lys Asp Tyr Val Glu Ile Asn	395	400	405
Gly Glu Lys Tyr	Cys Gly Glu Arg Ser Gln Phe Val Val Thr Ser	410	415	420
Asn Ser Asn Lys	Ile Thr Val Arg Phe His Ser Asp Gln Ser Tyr	425	430	435
Thr Asp Thr Gly	Phe Leu Ala Glu Tyr Leu Ser Tyr Asp Ser Ser	440	445	450
Asp Pro Cys Pro	Gly Gln Phe Thr Cys Arg Thr Gly Arg Cys Ile	455	460	465



Arg Lys Glu Leu Arg Cys Asp Gly Trp Ala Asp Cys Thr Asp His	470	475	480
Ser Asp Glu Leu Asn Cys Ser Cys Asp Ala Gly His Gln Phe Thr	485	490	495
Cys Lys Asn Lys Phe Cys Lys Pro Leu Phe Trp Val Cys Asp Ser	500	505	510
Val Asn Asp Cys Gly Asp Asn Ser Asp Glu Gln Gly Cys Ser Cys	515	520	525
Pro Ala Gln Thr Phe Arg Cys Ser Asn Gly Lys Cys Leu Ser Lys	530	535	540
Ser Gln Gln Cys Asn Gly Lys Asp Asp Cys Gly Asp Gly Ser Asp	545	550	555
Glu Ala Ser Cys Pro Lys Val Asn Val Val Thr Cys Thr Lys His	560	565	570
Thr Tyr Arg Cys Leu Asn Gly Leu Cys Leu Ser Lys Gly Asn Pro	575	580	585
Glu Cys Asp Gly Lys Glu Asp Cys Ser Asp Gly Ser Asp Glu Lys	590	595	600
Asp Cys Asp Cys Gly Leu Arg Ser Phe Thr Arg Gln Ala Arg Val	605	610	615
Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val	620	625	630
Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu	635	640	645
Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp	650	655	660
Asp Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe	665	670	675
Leu Gly Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln	680	685	690
Glu Arg Arg Leu Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp	695	700	705
Phe Thr Phe Asp Tyr Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro	710	715	720
Ala Glu Tyr Ser Ser Met Val Arg Pro Ile Cys Leu Pro Asp Ala	725	730	735
Ser His Val Phe Pro Ala Gly Lys Ala Ile Trp Val Thr Gly Trp	740	745	750

Gly His Thr Gln Tyr Gly Gly Thr Gly Ala Leu Ile Leu Gln Lys		
	755	760 765
Gly Glu Ile Arg Val Ile Asn Gln Thr Thr Cys Glu Asn Leu Leu		
	770	775 780
Pro Gln Gln Ile Thr Pro Arg Met Met Cys Val Gly Phe Leu Ser		
	785	790 795
Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro Leu Ser		
	800	805 810
Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly Val Val Ser		
	815	820 825
Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val Tyr Thr		
	830	835 840
Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr Gly Val		
	845	850 855

&lt;210&gt; 3

&lt;211&gt; 256

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of hepsin (Heps)  
homologous to similar domain in TADG-15

&lt;400&gt; 3

Arg Ile Val Gly Gly Arg Asp Thr Ser Leu Gly Arg Trp Pro Trp		
	5	10 15
Gln Val Ser Leu Arg Tyr Asp Gly Ala His Leu Cys Gly Gly Ser		
	20	25 30
Leu Leu Ser Gly Asp Trp Val Leu Thr Ala Ala His Cys Phe Pro		
	35	40 45
Glu Arg Asn Arg Val Leu Ser Arg Trp Arg Val Phe Ala Gly Ala		
	50	55 60
Val Ala Gln Ala Ser Pro His Gly Leu Gln Leu Gly Val Gln Ala		
	65	70 75
Val Val Tyr His Gly Gly Tyr Leu Pro Phe Arg Asp Pro Asn Ser		
	80	85 90
Glu Glu Asn Ser Asn Asp Ile Ala Leu Val His Leu Ser Ser Pro		
	95	100 105

Leu	Pro	Leu	Thr	Glu	Tyr	Ile	Gln	Pro	Val	Cys	Leu	Pro	Ala	Ala
				110					115					120
Gly	Gln	Ala	Leu	Val	Asp	Gly	Lys	Ile	Cys	Thr	Val	Thr	Gly	Trp
				125					130					135
Gly	Asn	Thr	Gln	Tyr	Tyr	Gly	Gln	Gln	Ala	Gly	Val	Leu	Gln	Glu
				140					145					150
Ala	Arg	Val	Pro	Ile	Ile	Ser	Asn	Asp	Val	Cys	Asn	Gly	Ala	Asp
				155					160					165
Phe	Tyr	Gly	Asn	Gln	Ile	Lys	Pro	Lys	Met	Phe	Cys	Ala	Gly	Tyr
				170					175					180
Pro	Glu	Gly	Gly	Ile	Asp	Ala	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro
				185					190					195
Phe	Val	Cys	Glu	Asp	Ser	Ile	Ser	Arg	Thr	Pro	Arg	Trp	Arg	Leu
				200					205					210
Cys	Gly	Ile	Val	Ser	Trp	Gly	Thr	Gly	Cys	Ala	Leu	Ala	Gln	Lys
				215					220					225
Pro	Gly	Val	Tyr	Thr	Lys	Val	Ser	Asp	Phe	Arg	Glu	Trp	Ile	Phe
				230					235					240
Gln	Ala	Ile	Lys	Thr	His	Ser	Glu	Ala	Ser	Gly	Met	Val	Thr	Gln
				245					250					255

Leu

<210>	4
<211>	225
<212>	PRT
<213>	Unknown
<220>	
<221>	DOMAIN
<223>	Serine protease catalytic domain of Scce homologous to similar domain in TADG-15.
<400>	4

Lys	Ile	Ile	Asp	Gly	Ala	Pro	Cys	Ala	Arg	Gly	Ser	His	Pro	Trp
				5					10					15
Gln	Val	Ala	Leu	Leu	Ser	Gly	Asn	Gln	Leu	His	Cys	Gly	Gly	Val
				20					25					30
Leu	Val	Asn	Glu	Arg	Trp	Val	Leu	Thr	Ala	Ala	His	Cys	Lys	Met
				35					40					45

```

Asn Glu Tyr Thr Val His Leu Gly Ser Asp Thr Leu Gly Asp Arg
      50                      55                      60
Arg Ala Gln Arg Ile Lys Ala Ser Lys Ser Phe Arg His Pro Gly
      65                      70                      75
Tyr Ser Thr Gln Thr His Val Asn Asp Leu Met Leu Val Lys Leu
      80                      85                      90
Asn Ser Gln Ala Arg Leu Ser Ser Met Val Lys Lys Val Arg Leu
      95                      100                     105
Pro Ser Arg Cys Glu Pro Pro Gly Thr Thr Cys Thr Val Ser Gly
      110                     115                     120
Trp Gly Thr Thr Thr Ser Pro Asp Val Thr Phe Pro Ser Asp Leu
      125                     130                     135
Met Cys Val Asp Val Lys Leu Ile Ser Pro Gln Asp Cys Thr Lys
      140                     145                     150
Val Tyr Lys Asp Leu Leu Glu Asn Ser Met Leu Cys Ala Gly Ile
      155                     160                     165
Pro Asp Ser Lys Lys Asn Ala Cys Asn Gly Asp Ser Gly Gly Pro
      170                     175                     180
Leu Val Cys Arg Gly Thr Leu Gln Gly Leu Val Ser Trp Gly Thr
      185                     190                     195
Phe Pro Cys Gly Gln Pro Asn Asp Pro Gly Val Tyr Thr Gln Val
      200                     205                     210
Cys Lys Phe Thr Lys Trp Ile Asn Asp Thr Met Lys Lys His Arg
      215                     220                     225

```

&lt;210&gt; 5

&lt;211&gt; 225

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of trypsin  
(Try) homologous to similar domain in TADG-15.

&lt;400&gt; 5

```

Lys Ile Val Gly Gly Tyr Asn Cys Glu Glu Asn Ser Val Pro Tyr
      5                      10                      15
Gln Val Ser Leu Asn Ser Gly Tyr His Phe Cys Gly Gly Ser Leu
      20                      25                      30

```

```

Ile Asn Glu Gln Trp Val Val Ser Ala Gly His Cys Tyr Lys Ser
      35                      40                      45
Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Glu Val Leu Glu
      50                      55                      60
Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro
      65                      70                      75
Gln Tyr Asp Arg Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys
      80                      85                      90
Leu Ser Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser
      95                      100                     105
Leu Pro Thr Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser
      110                     115                     120
Gly Trp Gly Asn Thr Ala Ser Ser Gly Ala Asp Tyr Pro Asp Glu
      125                     130                     135
Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Ala Lys Cys Glu
      140                     145                     150
Ala Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly
      155                     160                     165
Phe Leu Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly
      170                     175                     180
Pro Val Val Cys Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly
      185                     190                     195
Asp Gly Cys Ala Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val
      200                     205                     210
Tyr Asn Tyr Val Lys Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser
      215                     220                     225

```

&lt;210&gt; 6

&lt;211&gt; 231

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of chymotrypsin  
(Chymb) homologous to similar domain in TADG-15.

&lt;400&gt; 6

```

Arg Ile Val Asn Gly Glu Asp Ala Val Pro Gly Ser Trp Pro Trp
      5                      10                      15

```

Gln Val Ser Leu Gln Asp Lys Thr Gly Phe His Phe Cys Gly Gly	20	25	30
Ser Leu Ile Ser Glu Asp Trp Val Val Thr Ala Ala His Cys Gly	35	40	45
Val Arg Thr Ser Asp Val Val Val Ala Gly Glu Phe Asp Gln Gly	50	55	60
Ser Asp Glu Glu Asn Ile Gln Val Leu Lys Ile Ala Lys Val Phe	65	70	75
Lys Asn Pro Lys Phe Ser Ile Leu Thr Val Asn Asn Asp Ile Thr	80	85	90
Leu Leu Lys Leu Ala Thr Pro Ala Arg Phe Ser Gln Thr Val Ser	95	100	105
Ala Val Cys Leu Pro Ser Ala Asp Asp Asp Phe Pro Ala Gly Thr	110	115	120
Leu Cys Ala Thr Thr Gly Trp Gly Lys Thr Lys Tyr Asn Ala Asn	125	130	135
Lys Thr Pro Asp Lys Leu Gln Gln Ala Ala Leu Pro Leu Leu Ser	140	145	150
Asn Ala Glu Cys Lys Lys Ser Trp Gly Arg Arg Ile Thr Asp Val	155	160	165
Met Ile Cys Ala Gly Ala Ser Gly Val Ser Ser Cys Met Gly Asp	170	175	180
Ser Gly Gly Pro Leu Val Cys Gln Lys Asp Gly Ala Trp Thr Leu	185	190	195
Val Gly Ile Val Ser Trp Gly Ser Asp Thr Cys Ser Thr Ser Ser	200	205	210
Pro Gly Val Tyr Ala Arg Val Thr Lys Leu Ile Pro Trp Val Gln	215	220	225
Lys Ile Leu Ala Ala Asn	230		

&lt;210&gt; 7

&lt;211&gt; 255

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of factor 7  
(Fac7) homologous to similar domain in TADG-15.

&lt;400&gt; 7

Arg Ile Val Gly Gly Lys Val Cys Pro Lys Gly Glu Cys Pro Trp		
	5	10 15
Gln Val Leu Leu Leu Val Asn Gly Ala Gln Leu Cys Gly Gly Thr		
	20	25 30
Leu Ile Asn Thr Ile Trp Val Val Ser Ala Ala His Cys Phe Asp		
	35	40 45
Lys Ile Lys Asn Trp Arg Asn Leu Ile Ala Val Leu Gly Glu His		
	50	55 60
Asp Leu Ser Glu His Asp Gly Asp Glu Gln Ser Arg Arg Val Ala		
	65	70 75
Gln Val Ile Ile Pro Ser Thr Tyr Val Pro Gly Thr Thr Asn His		
	80	85 90
Asp Ile Ala Leu Leu Arg Leu His Gln Pro Val Val Leu Thr Asp		
	95	100 105
His Val Val Pro Leu Cys Leu Pro Glu Arg Thr Phe Ser Glu Arg		
	110	115 120
Thr Leu Ala Phe Val Arg Phe Ser Leu Val Ser Gly Trp Gly Gln		
	125	130 135
Leu Leu Asp Arg Gly Ala Thr Ala Leu Glu Leu Met Val Leu Asn		
	140	145 150
Val Pro Arg Leu Met Thr Gln Asp Cys Leu Gln Gln Ser Arg Lys		
	155	160 165
Val Gly Asp Ser Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala Gly		
	170	175 180
Tyr Ser Asp Gly Ser Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly		
	185	190 195
Pro His Ala Thr His Tyr Arg Gly Thr Trp Tyr Leu Thr Gly Ile		
	200	205 210
Val Ser Trp Gly Gln Gly Cys Ala Thr Val Gly His Phe Gly Val		
	215	220 225
Tyr Thr Arg Val Ser Gln Tyr Ile Glu Trp Leu Gln Lys Leu Met		
	230	235 240
Arg Ser Glu Pro Arg Pro Gly Val Leu Leu Arg Ala Pro Phe Pro		
	245	250 255

&lt;210&gt; 8

&lt;211&gt; 253

<212> PRT  
 <213> Unknown  
 <220>  
 <221> DOMAIN  
 <223> Serine protease catalytic domain of tissue plasminogen activator (Tpa) homologous to similar domain in TADG-15.  
 <400> 8

Arg	Ile	Lys	Gly	Gly	Leu	Phe	Ala	Asp	Ile	Ala	Ser	His	Pro	Trp	5	10	15
Gln	Ala	Ala	Ile	Phe	Ala	Lys	His	Arg	Arg	Ser	Pro	Gly	Glu	Arg	20	25	30
Phe	Leu	Cys	Gly	Gly	Ile	Leu	Ile	Ser	Ser	Cys	Trp	Ile	Leu	Ser	35	40	45
Ala	Ala	His	Cys	Phe	Gln	Glu	Arg	Phe	Pro	Pro	His	His	Leu	Thr	50	55	60
Val	Ile	Leu	Gly	Arg	Thr	Tyr	Arg	Val	Val	Pro	Gly	Glu	Glu	Glu	65	70	75
Gln	Lys	Phe	Glu	Val	Glu	Lys	Tyr	Ile	Val	His	Lys	Glu	Phe	Asp	80	85	90
Asp	Asp	Thr	Tyr	Asp	Asn	Asp	Ile	Ala	Leu	Leu	Gln	Leu	Lys	Ser	95	100	105
Asp	Ser	Ser	Arg	Cys	Ala	Gln	Glu	Ser	Ser	Val	Val	Arg	Thr	Val	110	115	120
Cys	Leu	Pro	Pro	Ala	Asp	Leu	Gln	Leu	Pro	Asp	Trp	Thr	Glu	Cys	125	130	135
Glu	Leu	Ser	Gly	Tyr	Gly	Lys	His	Glu	Ala	Leu	Ser	Pro	Phe	Tyr	140	145	150
Ser	Glu	Arg	Leu	Lys	Glu	Ala	His	Val	Arg	Leu	Tyr	Pro	Ser	Ser	155	160	165
Arg	Cys	Thr	Ser	Gln	His	Leu	Leu	Asn	Arg	Thr	Val	Thr	Asp	Asn	170	175	180
Met	Leu	Cys	Ala	Gly	Asp	Thr	Arg	Ser	Gly	Gly	Pro	Gln	Ala	Asn	185	190	195
Leu	His	Asp	Ala	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val	Cys	200	205	210



Leu Asn Asp Gly Arg Met Thr Leu Val Gly Ile Ile Ser Trp Gly  
 215 220 225  
 Leu Gly Cys Gly Gln Lys Asp Val Pro Gly Val Tyr Thr Lys Val  
 230 235 240  
 Thr Asn Tyr Leu Asp Trp Ile Arg Asp Asn Met Arg Pro  
 245 250

<210> 9  
 <211> 2900  
 <212> DNA  
 <213> *Homo sapiens*  
 <220>  
 <223> SNC19 mRNA sequence (U20428)  
 <400> 9

cgctgggtgg tgctggcagc cgtgctgata ggcctcctct tggctcttgc 50  
 ggggatcggc ttcttggtgt ggcatttgca gtaccgggac gtgcgtgtcc 100  
 agaaggtctt caatggctac atgaggatca caaatgagaa ttttgtggat 150  
 gcctacgaga actccaactc cactgagttt gtaagcctgg ccagcaagg 200  
 gaaggacgcg ctgaagctgc tgtacagcgg agtcccatc ctgggcccct 250  
 accacaagga gtcggctgtg acggccttca gcgagggcag cgtcatcgcc 300  
 tactactggt ctgagttcag catcccgag cacctggttg aggaggccga 350  
 gcgcgtcatg gccaggagcg cgtagtcatg ctgccccgc gggcgcgctc 400  
 cctgaagtcc tttgtggtca cctcagtggg ggctttcccc acggactcca 450  
 aaacagtaca gaggaccag gacaacagct gcagctttgg cctgcacgcc 500  
 gcggtgtgga gctgatgcgc ttcaccagc cggtttccct gacagcccct 550  
 accccgctca tgcccgtgc cagtgggctg cggggacgcg acgcagtgc 600  
 gagctactcg agctgactcg cagcttgact gcgcctcgac gagcgcgga 650  
 gcgacctggt gacgtgtaca acacctgag ccccatggag cccacgcct 700  
 ggtgagtgtg tggcacctac cctccctcct acaacctgac cttccactcc 750  
 ctcccacgaa cgtcctgctc atcacactga taaccaacac tgacgcggca 800  
 tcccggcttt gaggccacct tcttcagct gcctaggatg agcagctgtg 850  
 gaggccgett acgtaaagcc caggggacat tcaacagccc ctactacca 900  
 ggccactacc caccacacat tgactgcaca tggaaaattg aggtgcccac 950  
 caaccagcat gtgaaggtgc gttcaaatt cttctacctg ctggagcccc 1000  
 gcgtgcctgc gggcacctgc cccaaggact acgtggagat caatggggag 1050  
 aaatactgcg gagagaggtc ccagttcgtc gtcaccagca acagcaacaa 1100  
 gatcacagtt cgcttcact cagatcagtc ctacaccgac accggcttct 1150  
 tagctgaata cctctctac gactccagt acccatgccc ggggcagttc 1200  
 acgtgccgca cggggcggtg tatccggaag gagctgcgct gtgatggctg 1250

ggcgactgca cccaccacag cgatgagctc aactgcagtt gcgacgcgg 1300  
ccaccagttc acgtgcaaga gcaagttctg caagctcttc tgggtctgcg 1350  
acagtgtgaa cgagtgcgga gacaacagcg acgagcaggg ttgcatttgt 1400  
ccggacccag accttcaggt gttccaatgg gaagtgcctc tcgaaaagcc 1450  
agcagtgcaa tgggaaggac gactgtgggg acgggtccga cgaggcctcc 1500  
tgccccaagg tgaacgtcgt cacttgtacc aaacacacct accgctgcct 1550  
caatgggctc tgcttgagca agggcaacct tgagtgtgac gggaaggagg 1600  
actgtagcga cggctcagat gagaaggact gcgactgtgg gctgcggtca 1650  
ttcacgagac aggctcgtgt tgttgggggc acggatgcgg atgagggcga 1700  
gtggccctgg caggttaagcc tgcattctct gggccagggc cacatctgcg 1750  
gtgcttccct catctctccc aactggctgg tctctgccgc aacttgctac 1800  
atcgatgaca gaggattcag gtactcagac cccacgcagg acggccttcc 1850  
tgggcttgca cgaccagagc cagcgcaggc cctgggggtgc aggagcgcag 1900  
gctcaagcgc atcatctccc accccttctt caatgacttc accttcgact 1950  
atgacatcgc gctgctggag ctggagaaac cggcagagta cagctocatg 2000  
gtgcggccca tctgcctgcc ggacgcctgc catgtcttcc ctgccggcaa 2050  
ggccatctgg gtcacgggct ggggacacac ccagtatgga ggcaactggc 2100  
cgctgatcct gcaaaagggt gagatccgcg tcatcaacca gaccacctgc 2150  
gagaacctcc tgccgcagca gatcacgccg cgcattgatgt gcgtgggctt 2200  
cctcagcggc ggcggtggact cctgccaggg tgattccggg ggacctctgt 2250  
ccagcgtgga ggcggtatgg cggatcttcc aggcgggtgt ggtgagctgg 2300  
ggagacgctg cgctcagagg aacaagccag gcgtgtacac aaggctccct 2350  
ctgtttcggg aatggatcaa agagaacact ggggtatagg ggccggggcc 2400  
acccaatgt gtacacctgc ggggccaccc atcgtccacc ccagtgtgca 2450  
cgctgcagg ctggagactc gcgcaccgtg acctgcacca gcgccccaga 2500  
acatacactg tgaactcatc tccaggctca aatctgctag aaaacctctc 2550  
gcttccctcag cctccaaagt ggagctggga gggtagaagg ggaggaacac 2600  
tgggtggttct actgacccaa ctggggcaag gtttgaagca cagctccggc 2650  
agcccaagtg ggcgaggacg cgtttgtgca tactgccctg ctctatacac 2700  
ggaagacctg gatctctagt gagtgtgact gccggatctg gctgtggtcc 2750  
ttggccacgc ttcttgagga agcccaggct cggaggaccc tggaaaacag 2800  
acgggtctga gactgaaaat ggtttaccag ctcccagggtg acttcagtgt 2850  
gtgtattgtg taaatgagta aaacatttta tttcttttta aaaaaaaaaa 2900

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<221> primer  
<223> Forward primer for analysis of overexpression  
of TADG-15 mRNA by quantitative PCR.  
<400> 10  
atgacagagg attcaggtac 20  
<210> 11  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
<220>  
<221> primer  
<223> Reverse primer for analysis of overexpression  
of TADG-15 mRNA by quantitative PCR.  
<400> 11  
gaaggtgaag tcattgaaga 20  
<210> 12  
<211> 17  
<212> DNA  
<213> Artificial Sequence  
<220>  
<221> primer  
<223> Forward primer for analysis of  $\beta$ -tubulin mRNA  
expression by quantitative PCR.  
<400> 12  
tgcattgaca acgaggc 17  
<210> 13  
<211> 17  
<212> DNA  
<213> Artificial Sequence  
<220>  
<221> primer  
<223> Reverse primer for analysis of  $\alpha$ -tubulin mRNA  
expression by quantitative PCR.  
<400> 13  
ctgtcttgac attgttg 17

<210> 14  
 <211> 242  
 <212> PRT  
 <213> *Homo sapiens*  
 <220>  
 <221> DOMAIN  
 <223> Serine protease catalytic domain of TADG-15.  
 <400> 14

Arg Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp	5	10	15
Gln Val Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala	20	25	30
Ser Leu Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr	35	40	45
Ile Asp Asp Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr	50	55	60
Ala Phe Leu Gly Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly	65	70	75
Val Gln Glu Arg Arg Leu Lys Arg Ile Ile Ser His Pro Phe Phe	80	85	90
Asn Asp Phe Thr Phe Asp Tyr Asp Ile Ala Leu Leu Glu Leu Glu	95	100	105
Lys Pro Ala Glu Tyr Ser Ser Met Val Arg Pro Ile Cys Leu Pro	110	115	120
Asp Ala Ser His Val Phe Pro Ala Gly Lys Ala Ile Trp Val Thr	125	130	135
Gly Trp Gly His Thr Gln Tyr Gly Gly Thr Gly Ala Leu Ile Leu	140	145	150
Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr Thr Cys Glu Asn	155	160	165
Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met Cys Val Gly Phe	170	175	180
Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro	185	190	195
Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly Val	200	205	210
Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val	215	220	225

WO 99/42120

PCT/US99/03436

Tyr Thr Arg Leu Pro Leu Phe Arg Asp Trp Ile Lys Glu Asn Thr  
230 235 240

Gly Val

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/03436

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 530/324; 536/23.5; 435/320.1, 69.1, 6

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/324; 536/23.5; 435/320.1, 69.1, 6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DIALOG, APS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y,P	TANIMOTO, H. et al. Cloning and Expression of TADG-15, A Novel Serine Protease Expressed in Ovarian Cancer. Proceedings of the American Association for Cancer Research. March 1998, Vol. 39, page 648, especially page 648.	1-11
Y,P	O'BRIEN, T.J. et al. Cloning and Expression of TADG-15, A Novel Serine Protease Expressed in Ovarian Cancer" Tumor Biology. August 1998, Vol. 19, Supplement No. 2, pages 33, especially page 33.	1-11



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*U* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

26 APRIL 1999

Date of mailing of the international search report

19 MAY 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

YVONNE EYLER

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/03436

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 38/00; C07K 5/00, 7/00, 16/00, 17/00; C07H 21/04; C12N 15/00, 15/09, 15/63, 15/70, 15/74; C12P 21/06; C12Q  
1/68




FIG. 1



RIVGGRTSL GRMPWQVSL. ....RYDG.A HLCGGSLSG DWLTARCF PE....RNRV LSRWRVFAGA VACSPHGLC  
 RVGCTDADE GEMPWQVSL. ....HALGQS HICGASLISP NWLVSAHCY IDRGFRYSO PTQWTAFLGL HDQSQRSAFC  
 KIIDGAPCAR GSHPWQVAL. ....LSGNQL H.CGGYLVNE RNVLPAARHC. ....K MNEYTVHLGS OTLG...DR.F  
 KIVGGYNCEE NSVPTQVSL. ....NSGYHF ..CGGSLINE QWVSAHC. ....Y KSRIQVRIGE HNIEVLEG.F  
 RIVNGEDAVP GSWPWQVSL. ....QDKTGF HFCGSLISE DWVVTARHC. ....GV RTSQVVRAGE FQCGSDEE.F  
 RIVGGKVCCK GCEPWQVLL. ....LYMG.A QLCGGTLINT INWVSAHCF DKIKNMRNLI .....AVLGE HOLSEHDGDE  
 RIKGGGLFADI ASHPWQRAIF AKHRRSPGER FLCGGILISS CNILSAHCF QERPPPHHL. ....TVILGR .TYRVPFGEI

LGVQAVVYHG GYLPERDPNS EENSNDIALV HLSS.FLPLT EXIQPVCLPA ...AGQALVD GKICTVTGNG NTQYVGQQL  
 VQERRLKRIY SHPFENDETF D...YDIALL ELEK.PAEYS SNVRPFCLEP ...ASHVEFA GKAIWVTGNG HTQYGGTG.F  
 AQRIKASKSE RHFGYSTQT. ...HYNDLMLV KINS.QARLS SNVKKVRLES ...RCE..PP GTTCTVSGNG TTTSFDTTFI  
 EQFINAAKII RHFOYORAT. ...LNDIMLI KLSS.RAVIN ARVSTISLPT ...APP..AT GTKCLISGNG NFASSGADYI  
 IQVLKIAKVF KNPRESILT. ...VNNDITLL KLAT.PARFS QTVSAVCLES ...ADDEPA GTLCATTGNG KTKYMWANKTI  
 QSRRAQVII P...STYVP GTTNHDIALL RLHQ.FVULT DHVVPCLPE RTESEKTLAF VRPSLVSGNG QLLDRGATAI  
 EQKFEVEKYI VHKEFDDTY D...NDIALL QLKSSSRCA QESSVVRVTC LPFADQLPD NTECELSGYG KHEALSPEFY

\*

GVLOEARVEI ISNDVNGSAD FYGN..QIKP KMFAGYPEG G.....IDA CQGDSCGPFV CEDSISRTPR WRLCGIVSMI  
 LILQKGEIRV INQTCE..N LLPQ..QITP RMFCVGFLEG G.....VDS CQGDSCGPL. ...SSVERDGR IFQAGVVSMI  
 SGLMCVDVRL ISPDCTRV. .YKD..LLEN SMLCAGIPDS K.....KNA CMGDSGGPLV C.....R.... GTLQGLVSMI  
 DELOCLDAFV LSQAKCEAS. .YPG..KITS NMFCVGFLEG G.....KDG CQGDSCGPFV C.....N.... GQLQGVWSMI  
 EKLOQAALPL LSNAECKS. .WGR..RITD VMICAG..AS G.....VSE CMGDSGGPLV C.....QKDEA WTLNGIVSMI  
 ELMVLNVFRL MTQDCLQSR KVGDSFNITE YMFAGYSDG S.....KDS CKGDSGGP... ..HATHYRGT WYLTGIVSMI  
 ERLKEAHVRL YPSSRCTSQH LUNRT..VTD NMLCAGDTRS GGFQANLHDA CQGDSCGPFV CLM....DGR WTLVGLIISM

T.GCALAQKE	GVYTRVSDER	EWIFQAIKTH	SEASEMVTOL	--	(SEQ. ID NO: 3)	Heps
D.GCAQRNKE	GVYTRVLER	EWIKENTGV	-----	--	(SEQ. ID NO: 14)	Tadg 15
TFPCGQENDP	GVYTRVCKET	KWINDTMKKH	R-----	--	(SEQ. ID NO: 4)	SCCE
D.GCAQRNKE	GVYTRVINYV	KWIKNTIARN	S-----	--	(SEQ. ID NO: 5)	TRY
SDTCS.TSSE	GVYARVTKLI	PWQKILARN	-----	--	(SEQ. ID NO: 6)	Chymb
Q.GCATVGHF	GVYTRVSQYI	EQLQKLMRSE	PRFGVLLRAP	FP	(SEQ. ID NO: 7)	Fac 7
.LGGGQEDVE	GVYTRVTNVL	DMIRDMRE	-----	--	(SEQ. ID NO: 8)	Tpa

FIG. 2




FIG. 3




Figure 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8

○ : Conserved amino acids of catalytic triad H, D, S

Figure 9.



1	MCSDRARKGG	GGPKDFGACL	KYNSRHEKVM	GLEEGVEFLP	VNNVRKVEKR	1
51	GPGRVWVLAA	VLIGLLLVLL	GIGFLVWHLO	YRDVRVQKVF	NGYMRITNEN	2
101	FVDAYENSNS	TEFVSLASKV	KDALKLLYSG	VPFLGPIHKE	SAVTAFSEGS	
151	VIAYYWSEFS	IPQHLVEEAE	RVMAEERVVM	LPFRARSLKS	FWVTEVVAFP	
201	TDSKTVQORTQ	DNSCSFGLRA	RGVELMRFTT	PGFPDSPYPA	HARCQWALRG	
251	DADSVLSLTF	RSFDLASODE	RGSOLVTVM	TLSPMRPHAL	VQLCGTYFPS	
301	YNLTFFHSSQN	VLLITLITNT	ERRHFGFEAT	FEQLDRMSSC	GGRLRKAQGT	3
351	ENSPYYPGRY	PPNIDCTWNI	EVPNNOHVKV	SFKFFYLLEP	GVFAGTCPKD	
401	YVEINGEKYC	GERSQFVWTS	NSMKITVREH	SDQSYTDTGF	LAEYLSYDSS	
451	DECPGQETCR	TGRCIRKEIR	CDGWADCTDI	SDELNCSDOA	GHQETCKNKE	
501	CKPLEWVCDG	VNDGDNDEE	QGCSCPAQTF	RCENGKCLSK	SQQCNKDDC	4
551	GDGSDERASCP	KVNVVTCTKH	TYRCLNGLCL	SKCNPECDGK	EDCSDEDEK	
601	DDDCGLRSFT	RQARVVGSTD	ADEGEWFWQV	SIHALGQGHI	CGASLISPNW	
651	LVSAABCYID	DRGERYSOPT	QMTAFGLHD	QSORSAPGWQ	ERRLKRIISH	
701	PEFNDFTFDY	QHALLELEKP	AFYSSMVRPI	CLPDASHVVF	AGKAIWVTGW	5
751	GHTQYGGTGA	LILQKGEIRV	INQTTCEILL	PQGITPRMMC	VGFLSGGVDS	
801	CQGDGGPLS	SVEADGRIED	AGVVSQWGDGC	AQRNKEGVYT	RLPLERDWIK	
851	ENTGV	(SEQ. ID NO: 2)				

\* : Conserved cysteine residue

**NXT** : Possible N-linked glycosylation site

**SDE** : Conserved SDE motif

○ : Potential cleavage site

○ : Conserved amino acids of catalytic triad H, D, S

1. Cytoplasmic domain

2. Transmembrane domain


3. CUB repeat

4. Ligand-binding repeat (class A motif)  
of LDL receptor like domain

5. Serine protease

FIG. 10

11/13



1. Cytoplasmic domain
2. Transmembrane domain
3. Extracellular domain
- 4-5. CUB repeat
- 6-9. Ligand-binding repeat (class A motif) of LDL receptor like domain
10. Serine protease

FIG. 11

12/13

LOCUS H2000128 2000 bp RNA 221 01-05-1987

DEFINITION Homo SMC19 RNA sequence.

ACCESSION U06418

KEYWORDS

SOURCE human,

ORGANISM Homo sapiens

EMBRYONIC; mitochondrial eukaryotes; Vertebrata; Chordata; Vertebrates; Eutheria; Primates; Catarrhini; Hominoidea; Homo.

1 (bases 1 to 2000)

AUTHORS Zhang, G., Cai, X., Song, L., Cao, Q., Jiang, L. and Shi, J. Z.

TITLE SMC19 gene in Homo sapiens

JOURNAL Unpublished

REFERENCE 1 (bases 1 to 2000)

AUTHORS Zhang, G.

TITLE Direct Subcloning

JOURNAL Submitted (30-JUN-1985) Shu Xue, Jinder Institute, Shanghai Medical University, Nanxizhou, 20007, Peoples Republic of China



## SEQUENCE LISTING

- <110> O'Brien, Timothy J.  
Tanimoto, Hirotochi
- <120> TADG-15: An Extracellular Serine Protease  
Overexpressed in Breast and Ovarian Carcinomas
- <130> D6064PCT
- <140> PCT/US99/03436
- <141> 1999-02-18
- <150> US 09/627,337
- <151> 1998-02-20
- <160> 14
- <210> 1
- <211> 3147
- <212> INA
- <213> Homo sapiens
- <220>
- <223> cDNA sequence of TADG-15
- <400> 1

```

tcaagagccg cctcggygta ccatggggag ccatcgggcc cgaagggcg 50
gagggggccc gaaggaactc ggcgggggac tcaagtacaa ctcccgccac 100
gagaaagtga atggcttggg ggaaggcgtg gaggttcctgc cagtcacaaa 150
cgtcaagaag gtggaanaag atggcccggy gcgctgggtg gtgctggcag 200
ccgtgctgat cggcctcttc ttggctcttg tgggcacggg ctctctggtg 250
tggcatttgc agtacccgga cgtgcgtgtc cagaaggctc tcaatggcta 300
catgaggatc acaastgaga attttctgga tgcctacgag aactccaaact 350
ccactgagtt tgaagcctg gccagcaagg tgaaggacgc gctgaagctg 400
ctgtacagcg gagtcccat cctggggccc taccacaagg agtcggctgt 450
gacggccttc agcgaggcca ggcacatgc ctactactgg tctgagttca 500
gcaccccgca gaacctggty gaggaggccg agcgctcat ggcgaggag 550
cgcgtagtoa tgcctgccc gggggcgccg tccctgaagt cctttgtggt 600
caactcagtg gtggctttcc ccacggactc caaacagta cagaggcccc 650
aggacaacag ctgcagcttt ggccctgcac ccgcgggtgt ggagctgatg 700
cgcttcacca cgcgggctt cctgacagc cctaccccg cccatgccc 750
ctgcccagtg gccctgcgg gggacgccg ctcagtgtg agcctcacct 800
tccgcagctt tgaacttgcg tccctgcgac agcgcgccag cgaactggty 850
acggtgtaca acaccttgag ccccatggag cccacgccc tggtycagtt 900
gtgtggcacc taccctccct cctacaacct gaccttccac tctccacaga 950
acgtctgtct cctcacactg ataaccaaca ctgagcgggc gcaccccgcc 1000

```

tttagggcca ccttcttcca gctgcctagg atgagcagct gtggaggccg 1050  
 cttacgtaaa gcccaggga cattcaacag cccctactac ccaggccact 1100  
 acccacccaa cattgactgc acatggaaca ttgaggtgdc caccaccag 1150  
 catgtgaagg tgagcttcaa attcttctac ctgctggagc cggggtgdc 1200  
 tgggggcacc tgccccaagg actacgtgga gatcaatggg gagaaatact 1250  
 ggggagagag gtccagttc gtctcaccac gcaacagcaa caagatcaca 1300  
 gtctgcttcc actcagatca gtctcaccac gacaccggct ccttagctga 1350  
 atacctctcc taagactcca gtgacctatg cccggggcag ttcaagtgc 1400  
 gcacggggcg gtgtatccg aaggagctgc gctgtgatgg ctgggdcgac 1450  
 tgcccgacc adaggatga gctcaactgc agttgcgacg cccgccacca 1500  
 gttcacgtgc aagaacaagt tctgcaagcc cctcttctgg gtctgggaca 1550  
 gtgtgaacga ctggggagac aacagcgacg agcaggggtg cagttgtccg 1600  
 gccagacct tcaggtgttc caatgggaag tgcctctoga aaagccagca 1650  
 gtgcaatggg aaggacgact gtggggacgg gtccgacgag gcttctgdc 1700  
 ccaaggtgaa cgtcgtcact tgtaccaaac acacctaccg ctgctcact 1750  
 gggctctgct tgagcaaggy cacccttgag tgtgacggga aggaggactg 1800  
 tagcgacggc tcagatgaga aggactgcga ctgtgggctg cggctcattca 1850  
 cgagacaggg tegtgttgtt gggggcaccg atcgggatga gggcgagtgg 1900  
 ccttggcagg taagcctgca tgcctctggg cagggccaca tctgcggtgc 1950  
 ttccctcctc tctcccaact ggctggtctc tgccgcacac tgcctacatg 2000  
 atgacagagg attcaggtac tcagacccca cgcagtgagc ggcttctg 2050  
 ggcttgacg accagagcca ggcagcgcc cctggggtgc aggagcgag 2100  
 gttcaagcgc atcctctcc accctctctt caatgacttc accttogact 2150  
 atgacatcgc gctgctggag ctggagaaac cggcagagta cagctccatg 2200  
 gtgcggccca tctgctgdc ggacgctcc catgtcttcc ctgcccgcga 2250  
 ggcctcttgg gtcacgggt ggggacacac ccagtatgga ggcactggcg 2300  
 cgtgatcct gcaaaagggt gagatcccg tcatcaacca gaccacctgc 2350  
 gagaaactcc tgccgcagca gatcagcgg cgcagtatgt gctggggtt 2400  
 cctcaggggc ggcgtggact cctgdcagg tgattccgg ggaacctgt 2450  
 ccagcgtgga ggcggatgg cggatcttcc aggcgggtgt ggtgagctgg 2500  
 ggagacggct ggcgtcagag gaacaagcca ggcgtgtaca caaggctccc 2550  
 tctgttctgg gactggatca aagagaacac tgggttatag gggccggggc 2600  
 cacccaaatg tgtacaccty cggggccacc catcgtccac ccagtgctgc 2650  
 acgctgcag gctggagact ggaccgctga ctgcacagc gcccacagaa 2700  
 catcacctgt gaactcaatc ttcagggtct caaatctgcc tagaaaacct 2750  
 ctgcttctt cagcctccaa agtgagcty ggaggtagaa ggggaggaca 2800  
 ctggtggttc tactgaacca actggggcca aggttttgaa gatcagcct 2850  
 ccccgccag ccccaagcty ggcggaggcg cgtttgtgta tatctgctc 2900

```

ccctgtctgt aaggagcagc gggacggag cttcggagcc tcttcagtga 2950
aggtggtggg gctgcgggat ctgggtctgt gggccttgg gccacgctct 3000
tgagggaagcc caggtctgga ggaacctgga aaacagacgg gtctgagact 3050
gaaattgttt taccagctcc caggttggac ttcagtgtgt gtatttgtgt 3100
aaatgggtaa aacaatttat ttctttttta aaaaaaaaaa aaaaaaa 3147

```

```

<210>      2
<211>      855
<212>      PRT
<213>      Homo sapiens
<220>
<223>      Amino acid sequence of TADG-15 encoded by cDNA.
<400>      2

```

```

Met Gly Ser Asp Arg Ala Arg Lys Gly Gly Gly Gly Pro Lys Asp
      5                                10                    15
Phe Gly Ala Gly Leu Lys Tyr Asn Ser Arg His Glu Lys Val Asn
      20                                25                    30
Gly Leu Glu Glu Gly Val Glu Phe Leu Pro Val Asn Asn Val Lys
      35                                40                    45
Lys Val Glu Lys His Gly Pro Gly Arg Trp Val Val Leu Ala Ala
      50                                55                    60
Val Leu Ile Gly Leu Leu Leu Val Leu Leu Gly Ile Gly Phe Leu
      65                                70                    75
Val Trp His Leu Gln Tyr Arg Asp Val Arg Val Gln Lys Val Phe
      80                                85                    90
Asn Gly Tyr Met Arg Ile Thr Asn Glu Asn Phe Val Asp Ala Tyr
      95                                100                   105
Glu Asn Ser Asn Ser Thr Glu Phe Val Ser Leu Ala Ser Lys Val
      110                               115                   120
Lys Asp Ala Leu Lys Leu Leu Tyr Ser Gly Val Pro Phe Leu Gly
      125                               130                   135
Pro Tyr His Lys Glu Ser Ala Val Thr Ala Phe Ser Glu Gly Ser
      140                               145                   150
Val Ile Ala Tyr Tyr Trp Ser Glu Phe Ser Ile Pro Gln His Leu
      155                               160                   165
Val Glu Glu Ala Glu Arg Val Met Ala Glu Glu Arg Val Val Met
      170                               175                   180

```

Leu	Pro	Pro	Arg	Ala	Arg	Ser	Leu	Lys	Ser	Phe	Val	Val	Thr	Ser	185	190	195
Val	Val	Ala	Phe	Pro	Thr	Asp	Ser	Lys	Thr	Val	Gln	Arg	Thr	Gln	200	205	210
Asp	Asn	Ser	Cys	Ser	Phe	Gly	Leu	His	Ala	Arg	Gly	Val	Glu	Leu	215	220	225
Met	Arg	Phe	Thr	Thr	Pro	Gly	Phe	Pro	Asp	Ser	Pro	Tyr	Pro	Ala	230	235	240
His	Ala	Arg	Cys	Gln	Trp	Ala	Leu	Arg	Gly	Asp	Ala	Asp	Ser	Val	245	250	255
Leu	Ser	Leu	Thr	Phe	Arg	Ser	Phe	Asp	Leu	Ala	Ser	Cys	Asp	Glu	260	265	270
Arg	Gly	Ser	Asp	Leu	Val	Thr	Val	Tyr	Asn	Thr	Leu	Ser	Pro	Met	275	280	285
Glu	Pro	His	Ala	Leu	Val	Gln	Leu	Cys	Gly	Thr	Tyr	Pro	Pro	Ser	290	295	300
Tyr	Asn	Leu	Thr	Phe	His	Ser	Ser	Gln	Asn	Val	Leu	Leu	Ile	Thr	305	310	315
Leu	Ile	Thr	Asn	Thr	Glu	Arg	Arg	His	Pro	Gly	Phe	Glu	Ala	Thr	320	325	330
Phe	Phe	Gln	Leu	Pro	Arg	Met	Ser	Ser	Cys	Gly	Gly	Arg	Leu	Arg	335	340	345
Lys	Ala	Gln	Gly	Thr	Phe	Asn	Ser	Pro	Tyr	Tyr	Pro	Gly	His	Tyr	350	355	360
Pro	Pro	Asn	Ile	Asp	Cys	Thr	Trp	Asn	Ile	Glu	Val	Pro	Asn	Asn	365	370	375
Gln	His	Val	Lys	Val	Ser	Phe	Lys	Phe	Phe	Tyr	Leu	Leu	Glu	Pro	380	385	390
Gly	Val	Pro	Ala	Gly	Thr	Cys	Pro	Lys	Asp	Tyr	Val	Glu	Ile	Asn	395	400	405
Gly	Glu	Lys	Tyr	Cys	Gly	Glu	Arg	Ser	Gln	Phe	Val	Val	Thr	Ser	410	415	420
Asn	Ser	Asn	Lys	Ile	Thr	Val	Arg	Phe	His	Ser	Asp	Gln	Ser	Tyr	425	430	435
Thr	Asp	Thr	Gly	Phe	Leu	Ala	Gln	Tyr	Leu	Ser	Tyr	Asp	Ser	Ser	440	445	450
Asp	Pro	Cys	Pro	Gly	Gln	Phe	Thr	Cys	Arg	Thr	Gly	Arg	Cys	Ile	455	460	465



Arg Lys Glu Leu Arg Cys Asp Gly Trp Ala Asp Cys Thr Asp His	470	475	480
Ser Asp Glu Leu Asn Cys Ser Cys Asp Ala Gly His Gln Phe Thr	485	490	495
Cys Lys Asn Lys Phe Cys Lys Pro Leu Phe Trp Val Cys Asp Ser	500	505	510
Val Asn Asp Cys Gly Asp Asn Ser Asp Glu Gln Gly Cys Ser Cys	515	520	525
Pro Ala Gln Thr Phe Arg Cys Ser Asn Gly Lys Cys Leu Ser Lys	530	535	540
Ser Gln Gln Cys Asn Gly Lys Asp Asp Cys Gly Asp Gly Ser Asp	545	550	555
Glu Ala Ser Cys Pro Lys Val Asn Val Val Thr Cys Thr Lys His	560	565	570
Thr Tyr Arg Cys Leu Asn Gly Leu Cys Leu Ser Lys Gly Asn Pro	575	580	585
Glu Cys Asp Gly Lys Gln Asp Cys Ser Asp Gly Ser Asp Glu Lys	590	595	600
Asp Cys Asp Cys Gly Leu Arg Ser Phe Thr Arg Gln Ala Arg Val	605	610	615
Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp Gln Val	620	625	630
Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala Ser Leu	635	640	645
Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr Ile Asp	650	655	660
Asp Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr Ala Phe	665	670	675
Leu Gly Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly Val Gln	680	685	690
Glu Arg Arg Leu Lys Arg Ile Ile Ser His Pro Phe Phe Asn Asp	695	700	705
Phe Thr Phe Asp Tyr Asp Ile Ala Leu Leu Glu Leu Glu Lys Pro	710	715	720
Ala Glu Tyr Ser Ser Met Val Arg Pro Ile Cys Leu Pro Asp Ala	725	730	735
Ser His Val Phe Pro Ala Gly Lys Ala Ile Trp Val Thr Gly Trp	740	745	750

Gly	His	Thr	Gln	Tyr	Gly	Gly	Thr	Gly	Ala	Leu	Ile	Leu	Gln	Lys
					755				760					765
Gly	Glu	Ile	Arg	Val	Ile	Asn	Gln	Thr	Thr	Cys	Gln	Asn	Leu	Leu
					770				775					780
Pro	Gln	Gln	Ile	Thr	Pro	Arg	Met	Met	Cys	Val	Gly	Phe	Leu	Ser
					785				790					795
Gly	Gly	Val	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Ser
					800				805					810
Ser	Val	Glu	Ala	Asp	Gly	Arg	Ile	Phe	Gln	Ala	Gly	Val	Val	Ser
					815				820					825
Trp	Gly	Asp	Gly	Cys	Ala	Gln	Arg	Asn	Lys	Pro	Gly	Val	Tyr	Thr
					830				835					840
Arg	Leu	Pro	Leu	Phe	Arg	Asp	Trp	Ile	Lys	Gln	Asn	Thr	Gly	Val
					845				850					855

&lt;210&gt; 3

&lt;211&gt; 256

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;230&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of hepsin (Heps)  
homologous to similar domain in TADG-15

&lt;400&gt; 3

Arg	Ile	Val	Gly	Gly	Arg	Asp	Thr	Ser	Leu	Gly	Arg	Trp	Pro	Trp
									5			10		15
Gln	Val	Ser	Leu	Arg	Tyr	Asp	Gly	Ala	His	Leu	Cys	Gly	Gly	Ser
									20			25		30
Leu	Leu	Ser	Gly	Asp	Trp	Val	Leu	Thr	Ala	Ala	His	Cys	Phe	Pro
									35			40		45
Glu	Arg	Asn	Arg	Val	Leu	Ser	Arg	Trp	Arg	Val	Phe	Ala	Gly	Ala
									50			55		60
Val	Ala	Gln	Ala	Ser	Pro	His	Gly	Leu	Gln	Leu	Gly	Val	Gln	Ala
									65			70		75
Val	Val	Tyr	His	Gly	Gly	Tyr	Leu	Pro	Phe	Arg	Asp	Pro	Asn	Ser
									80			85		90
Glu	Glu	Asn	Ser	Asn	Asp	Ile	Ala	Leu	Val	His	Leu	Ser	Ser	Pro
									95			100		105

Leu Pro Leu Thr Glu Tyr Ile Gln Pro Val Cys Leu Pro Ala Ala	110	115	120
Gly Gln Ala Leu Val Asp Gly Lys Ile Cys Thr Val Thr Gly Trp	125	130	135
Gly Asn Thr Gln Tyr Tyr Gly Gln Gln Ala Gly Val Leu Gln Glu	140	145	150
Ala Arg Val Pro Ile Ile Ser Asn Asp Val Cys Asn Gly Ala Asp	155	160	165
Phe Tyr Gly Asn Gln Ile Lys Pro Lys Met Phe Cys Ala Gly Tyr	170	175	180
Pro Glu Gly Gly Ile Asp Ala Cys Gln Gly Asp Ser Gly Gly Pro	185	190	195
Phe Val Cys Glu Asp Ser Ile Ser Arg Thr Pro Arg Trp Arg Leu	200	205	210
Cys Gly Ile Val Ser Trp Gly Thr Gly Cys Ala Leu Ala Gln Lys	215	220	225
Pro Gly Val Tyr Thr Lys Val Ser Asp Phe Arg Glu Trp Ile Phe	230	235	240
Gln Ala Ile Lys Thr His Ser Glu Ala Ser Gly Met Val Thr Gln	245	250	255

Leu

<210>	4
<211>	225
<212>	PRT
<213>	Unknown
<220>	
<221>	DOMAIN
<223>	Serine protease catalytic domain of Scca homologous to similar domain in TADG-15.
<400>	4

Lys Ile Ile Asp Gly Ala Pro Cys Ala Arg Gly Ser His Pro Trp	5	10	15
Gln Val Ala Leu Leu Ser Gly Asn Gln Leu His Cys Gly Gly Val	20	25	30
Leu Val Asn Glu Arg Trp Val Leu Thr Ala Ala His Cys Lys Met	35	40	45

Asn	Glu	Tyr	Thr	Val	His	Leu	Gly	Ser	Asp	Thr	Leu	Gly	Asp	Arg
				50					55					60
Arg	Ala	Gln	Arg	Ile	Lys	Ala	Ser	Lys	Ser	Phe	Arg	His	Pro	Gly
				65					70					75
Tyr	Ser	Thr	Gln	Thr	His	Val	Asn	Asp	Leu	Met	Leu	Val	Lys	Leu
				80					85					90
Asn	Ser	Gln	Ala	Arg	Leu	Ser	Ser	Met	Val	Lys	Lys	Val	Arg	Leu
				95					100					105
Pro	Ser	Arg	Cys	Glu	Pro	Pro	Gly	Thr	Thr	Cys	Thr	Val	Ser	Gly
				110					115					120
Trp	Gly	Thr	Thr	Thr	Ser	Pro	Asp	Val	Thr	Phe	Pro	Ser	Asp	Leu
				125					130					135
Met	Cys	Val	Asp	Val	Lys	Leu	Ile	Ser	Pro	Gln	Asp	Cys	Thr	Lys
				140					145					150
Val	Tyr	Lys	Asp	Leu	Leu	Glu	Asn	Ser	Met	Leu	Cys	Ala	Gly	Ile
				155					160					165
Pro	Asp	Ser	Lys	Lys	Asn	Ala	Cys	Asn	Gly	Asp	Ser	Gly	Gly	Pro
				170					175					180
Leu	Val	Cys	Arg	Gly	Thr	Leu	Gln	Gly	Leu	Val	Ser	Trp	Gly	Thr
				185					190					195
Phe	Pro	Cys	Gly	Gln	Pro	Asn	Asp	Pro	Gly	Val	Tyr	Thr	Gln	Val
				200					205					210
Cys	Lys	Phe	Thr	Lys	Trp	Ile	Asn	Asp	Thr	Met	Lys	Lys	His	Arg
				215					220					225

&lt;210&gt; 5

&lt;211&gt; 225

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of trypsin  
(Try) homologous to similar domain in TADG-15.

&lt;400&gt; 5

Lys	Ile	Val	Gly	Gly	Tyr	Asn	Cys	Glu	Glu	Asn	Ser	Val	Pro	Tyr
				5					10					15
Gln	Val	Ser	Leu	Asn	Ser	Gly	Tyr	His	Phe	Cys	Gly	Gly	Ser	Leu
				20					25					30

Ile Asn Glu Gln Trp Val Val Ser Ala Gly His Cys Tyr Lys Ser	35	40	45
Arg Ile Gln Val Arg Leu Gly Glu His Asn Ile Gln Val Leu Glu	50	55	60
Gly Asn Glu Gln Phe Ile Asn Ala Ala Lys Ile Ile Arg His Pro	65	70	75
Gln Tyr Asp Arg Lys Thr Leu Asn Asn Asp Ile Met Leu Ile Lys	80	85	90
Leu Ser Ser Arg Ala Val Ile Asn Ala Arg Val Ser Thr Ile Ser	95	100	105
Leu Pro Thr Ala Pro Pro Ala Thr Gly Thr Lys Cys Leu Ile Ser	110	115	120
Gly Trp Gly Asn Thr Ala Ser Ser Gly Ala Asp Tyr Pro Asp Glu	125	130	135
Leu Gln Cys Leu Asp Ala Pro Val Leu Ser Gln Ala Lys Cys Glu	140	145	150
Ala Ser Tyr Pro Gly Lys Ile Thr Ser Asn Met Phe Cys Val Gly	155	160	165
Phe Leu Glu Gly Gly Lys Asp Ser Cys Gln Gly Asp Ser Gly Gly	170	175	180
Pro Val Val Cys Asn Gly Gln Leu Gln Gly Val Val Ser Trp Gly	185	190	195
Asp Gly Cys Ala Gln Lys Asn Lys Pro Gly Val Tyr Thr Lys Val	200	205	210
Tyr Asn Tyr Val Lys Trp Ile Lys Asn Thr Ile Ala Ala Asn Ser	215	220	225

&lt;210&gt; 6

&lt;211&gt; 231

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of chymotrypsin  
(Chymb) homologous to similar domain in TADG-15.

&lt;400&gt; 6

Arg Ile Val Asn Gly Glu Asp Ala Val Pro Gly Ser Trp Pro Trp	5	10	15
---	---	----	----

Gln	Val	Ser	Leu	Gln	Asp	Lys	Thr	Gly	Phe	His	Phe	Cys	Gly	Gly
				20					25					30
Ser	Leu	Ile	Ser	Glu	Asp	Trp	Val	Val	Thr	Ala	Ala	His	Cys	Gly
				35					40					45
Val	Arg	Thr	Ser	Asp	Val	Val	Val	Ala	Gly	Glu	Phe	Asp	Gln	Gly
				50					55					60
Ser	Asp	Glu	Glu	Asn	Ile	Gln	Val	Leu	Lys	Ile	Ala	Lys	Val	Phe
				65					70					75
Lys	Asn	Pro	Lys	Phe	Ser	Ile	Leu	Thr	Val	Asn	Asn	Asp	Ile	Thr
				80					85					90
Leu	Leu	Lys	Leu	Ala	Thr	Pro	Ala	Arg	Phe	Ser	Gln	Thr	Val	Ser
				95					100					105
Ala	Val	Cys	Leu	Pro	Ser	Ala	Asp	Asp	Asp	Phe	Pro	Ala	Gly	Thr
				110					115					120
Leu	Cys	Ala	Thr	Thr	Gly	Trp	Gly	Lys	Thr	Lys	Tyr	Asn	Ala	Asn
				125					130					135
Lys	Thr	Pro	Asp	Lys	Leu	Gln	Gln	Ala	Ala	Leu	Pro	Leu	Leu	Ser
				140					145					150
Asn	Ala	Glu	Cys	Lys	Lys	Ser	Trp	Gly	Arg	Arg	Ile	Thr	Asp	Val
				155					160					165
Met	Ile	Cys	Ala	Gly	Ala	Ser	Gly	Val	Ser	Ser	Cys	Met	Gly	Asp
				170					175					180
Ser	Gly	Gly	Pro	Leu	Val	Cys	Gln	Lys	Asp	Gly	Ala	Trp	Thr	Leu
				185					190					195
Val	Gly	Ile	Val	Ser	Trp	Gly	Ser	Asp	Thr	Cys	Ser	Thr	Ser	Ser
				200					205					210
Pro	Gly	Val	Tyr	Ala	Arg	Val	Thr	Lys	Leu	Ile	Pro	Trp	Val	Gln
				215					220					225
Lys	Ile	Leu	Ala	Ala	Asn									
				230										

&lt;210&gt; 7

&lt;211&gt; 255

&lt;212&gt; PRT

&lt;213&gt; Unknown

&lt;220&gt;

&lt;221&gt; DOMAIN

<223> Serine protease catalytic domain of factor 7  
 (Fac7) homologous to similar domain in TADG-15.

&lt;400&gt; 7

Arg	Ile	Val	Gly	Gly	Lys	Val	Cys	Pro	Lys	Gly	Glu	Cys	Pro	Trp
			5						10					15
Gln	Val	Leu	Leu	Leu	Val	Asn	Gly	Ala	Gln	Leu	Cys	Gly	Gly	Thr
			20						25					30
Leu	Ile	Asn	Thr	Ile	Trp	Val	Val	Ser	Ala	Ala	His	Cys	Phe	Asp
			35						40					45
Lys	Ile	Lys	Asn	Trp	Arg	Asn	Leu	Ile	Ala	Val	Leu	Gly	Glu	His
			50						55					60
Asp	Leu	Ser	Glu	His	Asp	Gly	Asp	Glu	Gln	Ser	Arg	Arg	Val	Ala
			65						70					75
Gln	Val	Ile	Ile	Pro	Ser	Thr	Tyr	Val	Pro	Gly	Thr	Thr	Asn	His
			80						85					90
Asp	Ile	Ala	Leu	Leu	Arg	Leu	His	Gln	Pro	Val	Val	Leu	Thr	Asp
			95						100					105
His	Val	Val	Pro	Leu	Cys	Leu	Pro	Glu	Arg	Thr	Phe	Ser	Glu	Arg
			110						115					120
Thr	Leu	Ala	Phe	Val	Arg	Phe	Ser	Leu	Val	Ser	Gly	Trp	Gly	Gln
			125						130					135
Leu	Leu	Asp	Arg	Gly	Ala	Thr	Ala	Leu	Glu	Leu	Met	Val	Leu	Asn
			140						145					150
Val	Pro	Arg	Leu	Met	Thr	Gln	Asp	Cys	Leu	Gln	Gln	Ser	Arg	Lys
			155						160					165
Val	Gly	Asp	Ser	Pro	Asn	Ile	Thr	Glu	Tyr	Met	Phe	Cys	Ala	Gly
			170						175					180
Tyr	Ser	Asp	Gly	Ser	Lys	Asp	Ser	Cys	Lys	Gly	Asp	Ser	Gly	Gly
			185						190					195
Pro	His	Ala	Thr	His	Tyr	Arg	Gly	Thr	Trp	Tyr	Leu	Thr	Gly	Ile
			200						205					210
Val	Ser	Trp	Gly	Gln	Gly	Cys	Ala	Thr	Val	Gly	His	Phe	Gly	Val
			215						220					225
Tyr	Thr	Arg	Val	Ser	Gln	Tyr	Ile	Glu	Trp	Leu	Gln	Lys	Leu	Met
			230						235					240
Arg	Ser	Glu	Pro	Arg	Pro	Gly	Val	Leu	Leu	Arg	Ala	Pro	Phe	Pro
			245						250					255

&lt;210&gt; 8

&lt;211&gt; 253

<212> PRT  
 <213> Unknown  
 <220>  
 <221> DOMAIN  
 <223> Serine protease catalytic domain of tissue plasminogen activator (Tpa) homologous to similar domain in TADG-15.

<400> 8

Arg	Ile	Lys	Gly	Gly	Leu	Phe	Ala	Asp	Ile	Ala	Ser	His	Pro	Trp
			5						10					15
Gln	Ala	Ala	Ile	Phe	Ala	Lys	His	Arg	Arg	Ser	Pro	Gly	Glu	Arg
			20						25					30
Phe	Leu	Cys	Gly	Gly	Ile	Leu	Ile	Ser	Ser	Cys	Trp	Ile	Leu	Ser
			35						40					45
Ala	Ala	His	Cys	Phe	Gln	Glu	Arg	Phe	Pro	Pro	His	His	Leu	Thr
			50						55					60
Val	Ile	Leu	Gly	Arg	Thr	Tyr	Arg	Val	Val	Pro	Gly	Glu	Glu	Glu
			65						70					75
Gln	Lys	Phe	Glu	Val	Glu	Lys	Tyr	Ile	Val	His	Lys	Glu	Phe	Asp
			80						85					90
Asp	Asp	Thr	Tyr	Asp	Asn	Asp	Ile	Ala	Leu	Leu	Gln	Leu	Lys	Ser
			95						100					105
Asp	Ser	Ser	Arg	Cys	Ala	Gln	Glu	Ser	Ser	Val	Val	Arg	Thr	Val
			110						115					120
Cys	Leu	Pro	Pro	Ala	Asp	Leu	Gln	Leu	Pro	Asp	Trp	Thr	Glu	Cys
			125						130					135
Glu	Leu	Ser	Gly	Tyr	Gly	Lys	His	Glu	Ala	Leu	Ser	Pro	Phe	Tyr
			140						145					150
Ser	Glu	Arg	Leu	Lys	Glu	Ala	His	Val	Arg	Leu	Tyr	Pro	Ser	Ser
			155						160					165
Arg	Cys	Thr	Ser	Gln	His	Leu	Leu	Asn	Arg	Thr	Val	Thr	Asp	Asn
			170						175					180
Met	Leu	Cys	Ala	Gly	Asp	Thr	Arg	Ser	Gly	Gly	Pro	Gln	Ala	Asn
			185						190					195
Leu	His	Asp	Ala	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val	Cys
			200						205					210



Leu Asn Asp Gly Arg Met Thr Leu Val Gly Ile Ile Ser Trp Gly  
 215 220 225  
 Leu Gly Cys Gly Gln Lys Asp Val Pro Gly Val Tyr Thr Lys Val  
 230 235 240  
 Thr Asn Tyr Leu Asp Trp Ile Arg Asp Asn Met Arg Pro  
 245 250

<210> 9  
 <211> 2900  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <223> SNC19 mRNA sequence (U20428)  
 <400> 9

cgctgggtgg tgetggcagc cgtgctgac ggctctctct tggctcttgc 50  
 ggggataggg ttcttggtgt ggcatttcca gtacggggac gtgctgtgac 100  
 agaaggtctt caatggctac atgaggatca caaatgagaa ttttgtggat 150  
 gctacagaga actccazctc cactgagttt gtaagcctgg ccagcaaggt 200  
 gaaggacgag ctgaagctgc tgtacagagg agtcccatte ctgggcccct 250  
 accacaagga gtgggtgtg agggccttca gogagggcag cgtcatcgcc 300  
 tactactggt ctgagttcag catcccgag caacttggtt agggggccga 350  
 ggcggtcatg gccaggagcg cgtagtcctg ctgccccgcg gggcgcgctc 400  
 cctgaagtcc tttgtggtca cctcagtggt ggccttcccc accgaactca 450  
 aaacagtaca gaggaccag gacaaacgct gcagcttttg cctgcacgac 500  
 ggggtgtgga gctgctgcgc ttcaccacgc cggcttccct gacagcccct 550  
 accccgctca tgcctgctgc cagtgggctg cggggagcgg acgcagtgc 600  
 gagctactcg agctgactcg cagcttgact ggcctctgac gaggcgggca 650  
 gogacctggt gacgtgtaca acacactgag ccccatggag cccacgct 700  
 ggtgagtgby tggacctac cttccctctt acaacctgac cttractcc 750  
 ctcccacgaa cgtctgctc atcacactga taaccaaac tgacgaggca 800  
 tcccggtttt gaggccacct tcttccagct gcctaggtatg agcagctgtg 850  
 gaggccgctt acctaaagcc caggggacat tcaacagccc ctactaccca 900  
 ggccactacc cacccaacat tgactgcaca tggaaaattg aggtgcccaa 950  
 caaccagcat gtgaaggtgc gcttcaaat cttctacctg ctggagcccc 1000  
 gcggtgctgc gggcaactgc cccaaggact acgtggagat caatggggag 1050  
 aaatactgag gagagaggtc ccagttogtc gtcaccagca acagcaacaa 1100  
 gatcacagtt cgtttccact cagatcagtc ctacaccgac accggcttct 1150  
 tagctgaata cctctctac gactccagtg aacctgccc ggggcagttc 1200  
 acgtgcccga cggggcggtg tatccggaag gagctgcgct gtgatggctg 1250

```

ggcgactgca ccgaccacag cgtatgagctc aactgcagtt gcgacgcggg 1300
ccaccagttc acgtgcaaga gcaggtttct caagctcttc tgggtctggg 1350
acagtgtgaa cgagtgcgga gacaacagcg acgagcaggg ttgcatttgt 1400
ccggacccag accttcaggt gtcccaatgg gaagtgcctc tcgaaagcc 1450
agcagtgcac tgggaaggac gactgtgggg acgggtccga cggagctctc 1500
tgcocaaagg tgaacgtcgt cacttgtacc aaacacacct accgctgctt 1550
caatgggctc tgcctgagca agggcaccgc cgagtgtgac gggaaaggag 1600
actgtagcga cggctcagat gagaaggact gcgactgtgg gatgagggtc 1650
ttcacgagac aggtctgtgt tgttgggggc acggatgcgg atgaggggc 1700
gtggccctgg caggtaagcc tgcctgctct gggtcagggc cacatctggg 1750
gtgcttccct catctctccc aactgggtgg tctctggcgc acactgtctc 1800
atcgatgaca gaggattcag gtactcagac cccacgcagg accgcttctc 1850
tgggcttgca ccaccagagc cagcgcaggg cctgggggtg aggagcgcag 1900
gtccaaagcg atcctctccc accccttctt caatgacttc accttcgact 1950
atgacatcgc gctgctggag ctggagaaac cggcagagta cagctccatg 2000
gtgggggccc tctgcttgcg ggacgcttgc catgtcttcc ctgcccggca 2050
ggcctctctg gtcacgggct ggggcacac ccagtatgga ggcactggcg 2100
cgctgatcct gcaaaagggt gagatccgcg tcatcaacca gacaccttgc 2150
gagaacctcc tgcgcagca gatcacgcgg cgcctgatgt gctggggctt 2200
cctcagcggc ggctgtgact cctgcacggg tgattccggg ggaacctgtt 2250
ccagcgtgga ggcggtatgg cggatcttcc aggcgggtgt ggtgagctgg 2300
ggagacgctg cgtcagagg aacaagccag gctgttacac aaggctccct 2350
ctgtttcggg aatggatcaa agacaacact ggggtatagg ggcggggg 2400
accgaaatgt gtacacctgc ggggcacccc atcgtccacc ccgctgtgca 2450
cgctgcagg ctggagactc ggcacccgtg acctgcacca ggcggccaga 2500
acatacactg tgaactcctc tccaggttca aatctgctag aaaacctctc 2550
gcttctctcag cctccaaagt ggagctggga ggttagaagg ggaggaaac 2600
tggtygttct actgacccaa ctggggcaag gtttgaagca cagctccggc 2650
agcccaagtg ggcgaggacg cgtttgtgca tactgcctg ctctatacac 2700
ggaagacctg gatctctagt gactgtgact ggcggatctg gctgtggtcc 2750
ttggccacgc ktcttgagga agcccaggct cggaggaccc tggaaaaacg 2800
acgggtctga gactgaaaat ggtttaccag ctcccaggtg acttcagtgt 2850
gtgtatttgt taaatgagta aaacatttta tttcttttta aaaaaaaa 2900

```

&lt;210&gt; 10

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<221> primer  
<223> Forward primer for analysis of overexpression  
of TADG-15 mRNA by quantitative PCR.  
<400> 10  
atgacagagg attcaggtac 20  
<210> 11  
<211> 20  
<212> DNA  
<213> Artificial Sequence  
<220>  
<221> primer  
<223> Reverse primer for analysis of overexpression  
of TADG-15 mRNA by quantitative PCR.  
<400> 11  
gaaggtgaag tcattgaaga 20  
<210> 12  
<211> 17  
<212> DNA  
<213> Artificial Sequence  
<220>  
<221> primer  
<223> Forward primer for analysis of  $\beta$ -tubulin mRNA  
expression by quantitative PCR.  
<400> 12  
tgcattgaca acgagggc 17  
<210> 13  
<211> 17  
<212> DNA  
<213> Artificial Sequence  
<220>  
<221> primer  
<223> Reverse primer for analysis of  $\beta$ -tubulin mRNA  
expression by quantitative PCR.  
<400> 12  
ctgtcttgac attgttcg 17

<210> 14  
 <211> 242  
 <212> PRT  
 <213> Homo sapiens  
 <220>  
 <221> DOMAIN  
 <223> Serine protease catalytic domain of TADG-15.  
 <400> 14

Arg Val Val Gly Gly Thr Asp Ala Asp Glu Gly Glu Trp Pro Trp		
	5	10 15
Gln Val Ser Leu His Ala Leu Gly Gln Gly His Ile Cys Gly Ala		
	20	25 30
Ser Leu Ile Ser Pro Asn Trp Leu Val Ser Ala Ala His Cys Tyr		
	35	40 45
Ile Asp Asp Arg Gly Phe Arg Tyr Ser Asp Pro Thr Gln Trp Thr		
	50	55 60
Ala Phe Leu Gly Leu His Asp Gln Ser Gln Arg Ser Ala Pro Gly		
	65	70 75
Val Gln Glu Arg Arg Leu Lys Arg Ile Ile Ser His Pro Phe Phe		
	80	85 90
Asn Asp Phe Thr Phe Asp Tyr Asp Ile Ala Leu Leu Glu Leu Glu		
	95	100 105
Lys Pro Ala Glu Tyr Ser Ser Met Val Arg Pro Ile Cys Leu Pro		
	110	115 120
Asp Ala Ser His Val Phe Pro Ala Gly Lys Ala Ile Trp Val Thr		
	125	130 135
Gly Trp Gly His Thr Gln Tyr Gly Gly Thr Gly Ala Leu Ile Leu		
	140	145 150
Gln Lys Gly Glu Ile Arg Val Ile Asn Gln Thr Thr Cys Glu Asn		
	155	160 165
Leu Leu Pro Gln Gln Ile Thr Pro Arg Met Met Cys Val Gly Phe		
	170	175 180
Leu Ser Gly Gly Val Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro		
	185	190 195
Leu Ser Ser Val Glu Ala Asp Gly Arg Ile Phe Gln Ala Gly Val		
	200	205 210
Val Ser Trp Gly Asp Gly Cys Ala Gln Arg Asn Lys Pro Gly Val		
	215	220 225